arXiv:1212.1378v1 [hep-th] 6 Dec 2012

LATTICE W-ALGEBRAS AND LOGARITHMIC CFTS

A.M. GAINUTDINOV, H. SALEUR AND I.YU. TIPUNIN

Abstract. This paper is part of an effort to gain further understanding of 2D Logarithmic Conformal Field Theories (LCFTs) by exploring their lattice regularizations. While all work so far has dealt with the Virasoro algebra (or the product $\mathrm{Vir}\otimes\overline{\mathrm{Vir}}$), the best known (although maybe not the most relevant physically) LCFTs in the continuum are characterized by a W-algebra symmetry, whose presence is powerful, but difficult to understand physically. We explore here the origin of this symmetry in the underlying lattice models. We consider $U_{\mathfrak{q}}s\ell(2)$ XXZ spin chains for $\mathfrak{q}$ a root of unity, and argue that the centralizer of the "small" quantum group $\overline{U}_{\mathfrak{q}}s\ell(2)$ goes over the W-algebra in the continuum limit. We justify this identification by representation theoretic arguments, and give, in particular, lattice versions of the W-algebra generators. In the case $\mathfrak{q}=i$, which corresponds to symplectic fermions at central charge $c=-2$, we provide a full analysis of the scaling limit of the lattice Virasoro and W generators, and show in details how the corresponding continuum Virasoro and W-algebras areobtained. Striking similarities between the lattice W algebra and the Onsager algebra are observed in this case.

1. Introduction

It is not surprising that some crucial algebraic objects should play a role both in lattice models and in conformal field theories. For instance, the Yang Baxter equation expresses factorizability of lattice interactions and determines integrable Boltzmann weights; it also expresses factorizability of braiding in the CFT, and is related with monodromy properties of conformal blocks.

Work of the last many years however has unraveled deeper and less expected connections. Among these was the surprising observation that order parameters (local height probabilities) are given by CFT branching functions [1], or that the representation theory of lattice algebras such as the Temperley–Lieb algebra is equivalent, in a certain sense (see [2, 3, 5] for more details), to the representation theory of the Virasoro algebra (see also [7, 8]). This has proven particularly useful in understanding logarithmic conformal field theories (LCFTs), a topic which has gained a lot of attention recently. While classifying and solving LCFTs from first principles seems exceedingly hard, the connection with lattice models allows one to rely on many advanced results of algebra - in particular on the representation theory of non semi-simple cellular algebras - to gain understanding of the types of indecomposable modules or fusion rules appearing in LCFTs of the most importance for applications [9].

It is crucial to understand the observations in [7, 3] more thoroughly, building on the intuitive notion that the algebra of local hamiltonians (e.g. nearest coupling for spin chains) should in some sense go over the algebra of local hamiltonians (the stress energy tensor) in the continuum limit, and for boundary theories, where only one chiral algebra is expected. However, showing precisely how say the Virasoro algebra emerges from the Temperley–Lieb algebra in the continuum limit remains a very hard exercise in general, except for the Ising model where the presence of an underlying Majorana fermion makes things quite easy [10].

It is also worth mentioning that among LCFTs the rational class with "large" chiral algebras like the triplet W-algebras [11, 4] has a finite number of primary fields (or isomorphism classes of simple modules) and is particularly accessible for rigorous algebraic analysis [13, 14], as well as for an explicit treatment of their Verlinde algebra aspects [15, 16] and hopefully mapping class group properties [17]. The simplest such LCFT occurs in the so called symplectic fermion theory at $c=-2$ [18, 11] where the chiral algebra is generated by a triplet of fermion bilinears of conformal weight $h=3$:

$$\begin{aligned} W^+ &= \partial\eta^+\eta^+, \\ W^0 &= \frac{1}{2}\left(\partial\eta^+\eta^- + \partial\eta^-\eta^+\right), \\ W^- &= \partial\eta^-\eta^- \end{aligned}$$

satisfying the non linear relations

$$\begin{aligned} W^\alpha(z)W^\beta(w) &= g^{\alpha\beta}\left(\frac{\frac{1}{2}}{(z-w)^6} - \frac{3}{2}\frac{T(w)}{(z-w)^4} - \frac{3}{4}\frac{\partial T(w)}{(z-w)^3}\right. \\ &\quad \left. +\frac{3}{8}\frac{\partial^2 T(w)}{(z-w)^2} - 2\frac{(T^2)(w)}{(z-w)^2} + \frac{1}{4}\frac{\partial^3 T(w)}{z-w} - \frac{\partial(T^2)(w)}{z-w}\right) \\ &\quad +5f_\gamma^{\alpha\beta}\left(\frac{W^\gamma(w)}{(z-w)^3} + \frac{1}{2}\frac{\partial W^\gamma(w)}{(z-w)^2} + \frac{1}{25}\frac{\partial^2 W^\gamma(w)}{z-w} + \frac{12}{25}\frac{(TW^\gamma)(w)}{z-w}\right) \end{aligned} \tag{1}$$

with $g^{+-}=g^{-+}=4$, $g^{00}=2$, $f_0^{+-}=-f_0^{-+}=2$, $f_+^{0+}=-f_+^{+0}=-f_-^{0-}=f_-^{-0}=1$ and other $g^{\alpha\beta}$ and $f_\gamma^{\alpha\beta}$ equal to 0.

Most studied generalizations concern the so called $(1,p)$ models (symplectic fermions being the $(1,2)$ model). Identifying lattice models which have such W-algebra symmetries remains, in our opinion and except for $p=2$, a challenge, even though some formal progress has been made in this direction in the last few years [19].

The study of the W-algebras is notoriously difficult, not only because of the high non linearity of the OPEs, but also because they do lack physical understanding. Motivated by the crucial role played by these algebras in the rational class of LCFTs, we are presenting here a fresh look at this question by identifying the lattice equivalents of the tripet W-algebras in $U_{\mathfrak{q}}s\ell(2)$ symmetric XXZ spin chains when $\mathfrak{q}$ is a root of unity. The idea of their definition is based on the full symmetry algebra (or *centralizer*) paradigm. So, the lattice equivalent of the Virasoro algebra could be defined using its centralizer in CFT [3, 25] –

the *full* quantum group $U_{\mathfrak{q}}s\ell(2)$ – as the algebra of local Hamiltonian densities commuting with the $U_{\mathfrak{q}}s\ell(2)$. Indeed, such an algebra on the finite spin-chain is the Temperley–Lieb (TL) algebra and it has close relations with the representation theory of the Virasoro algebra [2, 32]. Moreover, we will show in this paper that particular elements of the TL algebra in the XX spin-chain converge in the scaling limit to the Virasoro modes. To go further and define lattice algebras that would converge to the chiral tripet W-algebras, we use the (so called Kazhdan–Lusztig) duality [26, 12] between these chiral algebras and the *small* quantum group denoted by $\overline{U}_{\mathfrak{q}}s\ell(2)$. This duality essentially states[1] that the centralizer of the W-algebra in a particular Hilbert space of CFT states is the $\overline{U}_{\mathfrak{q}}s\ell(2)$. Therefore in this case, one could think about the lattice equivalent as the centralizer of the small quantum group on a finite spin-chain.

The paper is organized as follows. Sections 2 and 3 are devoted to the case $\mathfrak{q} = i$, that is the XX spin chain (with boundary terms), which is equivalent to a $g\ell(1|1)$ spin chain built on alternating representations. Using the earlier proposal in [10, 5], a lattice version of the Virasoro algebra is obtained by considering TL generators – recall that the TL algebra is the centralizer of the full quantum group acting on the spin chain. In section 2, we propose similarly a lattice version of the W-algebra obtained as the centralizer of the small quantum group $\overline{U}_{\mathfrak{q}}s\ell(2)$ acting on the spin chain. Fourier modes (particular linear combinations of the lattice W-algebra generators) which constitute "approximations" of the Virasoro or chiral W-algebra modes are then defined, and their commutators are studied in section 2 as well. Of course, for a finite system, these approximations do not form a closed algebra. Calculating further commutators leads to infinitely many such "approximations", and a complicated algebraic structure, which is studied in details. The scaling limit is then studied in section 3 where we are able, after the introduction of appropriate normal ordering, to obtain exactly, from the lattice and in the limit of large chains, the defining relations of the Virasoro and W algebras for symplectic fermions. In section 4, we generalize our analysis to other values of $\mathfrak{q}$. We define the lattice W algebra for the XXZ spin chain, and study its representation theory in details. While refraining from a full analysis of the continuum limit, we show that our results are fully consistent with what is expected in the $(p-1,p)$ and $(1,p)$ models of LCFTs. Section 5 contains some conclusions and pointers for future work. Five appendices contain supplementary materials like long definitions, proofs and bulky expressions.

2. The triplet W-algebra from XX spin-chain

In this section, we give an explicit definition of the lattice analogue of Kausch's triplet W-algebra in the case of XX spin-chains. The use of well-known fermionic formulation of the

[1]The equivalence between the categories of the triplet W-algebra modules and finite-dimensional $\overline{U}_{\mathfrak{q}}s\ell(2)$-modules was proven first in [12] for the $(1,2)$ models and in [27] for all $(1,p)$ cases. The functor realizing such an equivalence could be explicitly constructed using a bimodule over the mutual centralizers, the triplet W-algebra and the small quantum group.

model allows to introduce all modes of "W-currents" on the lattice. Their scaling limit and commutation relations are then studied in the next section.

2.1. **The XX model.** We consider the XX-hamiltonian of N one-half spins with an open boundary condition described by the "quantum-group symmetric" boundary term [23],

$$H = \frac{1}{2}\sum_{i=1}^{N-1}(\sigma_i^x\sigma_{i+1}^x + \sigma_i^y\sigma_{i+1}^y) - \frac{i}{2}(\sigma_1^z - \sigma_N^z), \tag{2}$$

where σ_i^x, σ_i^y and σ_i^z are usual Pauli matrices,

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{3}$$

2.2. **Lattice fermions.** It is useful in what follows to reformulate everything in terms of ordinary lattice fermions c_j and $c_j^\dagger$. We use the Jordan-Wigner transformation to get the lattice fermions

$$\begin{aligned} c_j^\dagger &= i^{j-1}\ i^{\sigma_1^z+\ldots+\sigma_{j-1}^z} \otimes \sigma_j^+, \\ c_j &= i^{-j+1}\ i^{-\sigma_1^z-\ldots-\sigma_{j-1}^z} \otimes \sigma_j^- \end{aligned} \tag{4}$$

with the anticommutation relations

$$\{c_j^\dagger, c_{j'}\} = \delta_{j,j'}. \tag{5}$$

This transformation gives the free fermion expression for the XX hamiltonian from (2):

$$H = -\sum_{j=1}^{N-1} e_j = \sum_{j=1}^{N-1}\big(c_j^\dagger c_{j+1} + c_{j+1}^\dagger c_j\big) - ic_1^\dagger c_1 + ic_N^\dagger c_N, \tag{6}$$

where we introduce the hamiltonian densities

$$e_j = c_j c_{j+1}^\dagger + c_{j+1}c_j^\dagger + i\big(c_j^\dagger c_j - c_{j+1}^\dagger c_{j+1}\big), \qquad 1 \le j \le N-1, \tag{7}$$

which satisfy defining relations for the Temperley–Lieb algebra with zero fugacity parameter:

$$\begin{aligned} e_i^2 &= 0, \\ e_i e_{i\pm1} e_i &= e_i, \\ e_i e_j &= e_j e_i, \quad |i-j| > 1. \end{aligned}$$

We denote the algebra generated by these e_j, with $1 \le j \le N-1$, and the identity as $\mathcal{TL}_{i,N}$.

2.3. **Quantum group results for $\mathfrak{q}=i$ case.** For applications to open XX spin-chains, we set in what follows $\mathfrak{q}\equiv i$. As a module over the quantum group $U_{\mathfrak{q}}s\ell(2)$, the spin chain $\mathcal{H}_N$ is a tensor product of two-dimensional irreducibe representations such that $\mathsf{E}\to\sigma^+$, $\mathsf{F}\to\sigma^-$, $\mathsf{K}\to\mathfrak{q}\sigma^z$, and $\mathsf{e}=\mathsf{f}=0$, see precise definitions of quantum groups at roots of unity in App. **A**. Using the $(N-1)$-folded comultiplications (145) together with the Jordan-Wigner transformation (4), we obtain the representation $\rho_{g\ell}: U_{\mathfrak{q}}s\ell(2)\to \mathrm{End}_{\mathbb{C}}(\mathcal{H}_N)$ (usual fermionic expressions)

$$\rho_{g\ell}(\mathsf{E})\equiv\Delta^{N-1}(\mathsf{E})=\sum_{j=1}^{N}\mathfrak{q}^j c_j^\dagger\mathsf{K},\qquad \rho_{g\ell}(\mathsf{F})\equiv\Delta^{N-1}(\mathsf{F})=\sum_{j=1}^{N}\mathfrak{q}^{j-1}c_j, \tag{8}$$

and

$$\begin{aligned}\rho_{g\ell}(\mathsf{e})\equiv\Delta^{N-1}(\mathsf{e})&=\sum_{1\le j_1<j_2\le N}(-1)^{j_1+j_2}\mathfrak{q}^{1-j_1-j_2}c_{j_1}^\dagger c_{j_2}^\dagger,\\ \rho_{g\ell}(\mathsf{f})\equiv\Delta^{N-1}(\mathsf{f})&=\sum_{1\le j_1<j_2\le N}\mathfrak{q}^{j_1+j_2-1}c_{j_1}c_{j_2}.\end{aligned} \tag{9}$$

We then can easily check that all the generators e_j of $\mathcal{TL}_{i,N}$ commutes with the quantum-group action [23]. Moreover, the quantum-group generators give the full symmetry of the XX model, or in more technical terms $U_{\mathfrak{q}}s\ell(2)$ at $\mathfrak{q}=i$ gives *the centralizer* of $\mathcal{TL}_{i,N}$. This is a particular case of a more general (and, of course, well-known) statement discussed in Sec. **4** in the context of XXZ representations.

2.4. **Lattice W-algebra: generators and their relations.** As was mentioned in the introduction, our paradigm in definition of lattice algebras (as well as chiral algebras in the scaling limit) is based on their centralizers which are various quantum groups. We will see in Sec. **3** that a properly defined scaling limit of $\mathcal{TL}_{i,N}$ gives all Virasoro modes L_n in symplectic fermions representation, a logarithmic CFT with the central charge $c=-2$. It was discussed in [3], see also [25], that the scaling limit of the $U_{\mathfrak{q}}s\ell(2)$ generators gives the centralizer of this Virasoro representation, or more formally, an equivalent "staircase" bimodule structure for the commuting actions of the Virasoro and $U_{\mathfrak{q}}s\ell(2)$ is obtained as a semi-infinite version of the finite ones for $\mathcal{TL}_{i,N}$ and $U_{\mathfrak{q}}s\ell(2)$ extracted from the XX spin-chains.

In the case of the chiral triplet W-algebra, it was proven in [12] that the centralizer of the W-algebra in the symplectic fermions is given by the small (or restricted) quantum group $\overline{U}_{\mathfrak{q}}s\ell(2)$ generated by the capital generators E and F, with the Cartan element K, only. Then, by an analogy with the Temperley–Lieb case, we define a lattice analogue of the Kausch's triplet W-algebra as an extension of the Temperley–Lieb algebra for $\mathfrak{q}=i$ by generators that respect the action of the restricted quantum group but break the $s\ell(2)$ symmetry generated by e and f. Saying more formally, we define *the lattice W-algebra* $\mathcal{W}_{i,N}$ as the algebra of all operators intertwining the action of the restricted quantum group $\overline{U}_{\mathfrak{q}}s\ell(2)$ on the tensor-product representation $\mathcal{H}_N$.

In order to find the additional generators extending $\mathcal{TL}_{i,N}$ to $\mathcal{W}_{i,N}$, we first recall a definition of the wall Brauer algebra in the context of alternating $g\ell(1|1)$ spin-chains introduced in [9].

2.4.1. *A relation with $g\ell(1|1)$ spin-chains and the wall Brauer algebra.* Up to some simple transformations, the XX spin chain can be reformulated as a $g\ell(1|1)$ spin chain with alternating fundamental representation and its dual. This is discussed in details for instance in the introductory part of [5] where it is shown, among other things, that $g\ell(1|1)$ symmetry is generated by the quantum-group generators E, F, and the $s\ell(2)$ Cartan element h via

$$\begin{aligned}\rho_{g\ell}(\mathsf{E}) &\equiv \Delta^{N-1}(\mathsf{E}) = F^\dagger_{(1)}\,\rho_{g\ell}(\mathsf{K}),\\ \rho_{g\ell}(\mathsf{F}) &\equiv \Delta^{N-1}(\mathsf{F}) = \mathsf{q}^{-1}F_{(1)},\\ \rho_{g\ell}(\mathsf{K}) &\equiv \Delta^{N-1}(\mathsf{K}) = (-1)^{\rho_{g\ell}(2\mathsf{h})},\end{aligned} \tag{10}$$

where $F^\dagger_{(1)}, F_{(1)}$ are the fermionic generators and $2\mathsf{h}$ the fermion number (the last generator of $g\ell(1|1)$ is trivially zero in the alternating product of fundamental and its dual).

We note now that the $g\ell(1|1)$ action commutes with more than the nearest-neighbor coupling – Temperley Lieb generators. It commutes in particular with permutations of next-nearest neighbor representations, which in mathematical terms corresponds to an action of the wall Brauer algebra [20] generated by e_j and WB_j. The wall Brauer generators are represented in our spin-chain by

$$WB_j = (c_j + c_{j+2})(c_j^\dagger + c_{j+2}^\dagger) - 1, \qquad j = 1, 2, \dots, N-2. \tag{11}$$

It was proven in [29] that this representation of the wall Brauer algebra gives the centralizer of the alternating $g\ell(1|1)$ tensor-space representation. We thus see that $WB_j \in \mathcal{W}_{i,N}$ because of the relations (10) which show the the $g\ell(1|1)$ representation contains the image of the restricted quantum group $\overline{U}_\mathsf{q} s\ell(2)$. Obviously, WB_j's do not commute with the $s\ell(2)$ generators e and f but still commute with the commutator $[\mathsf{e},\mathsf{f}] = 2\mathsf{h} = S^z$. In order to find the full centralizer of $\overline{U}_\mathsf{q} s\ell(2)$ with the Cartan K represented by the operator $(-1)^{2\mathsf{h}}$, we then introduce two more generators by

$$WB_j^+ = [\mathsf{e}, WB_j], \qquad WB_j^- = [\mathsf{f}, WB_j]. \tag{12}$$

These operators now break the S^z-symmetry (the $g\ell(1|1)$ generator which is not in $\overline{U}_\mathsf{q} s\ell(2)$) but respect the action of the Cartan generator K of $\overline{U}_\mathsf{q} s\ell(2)$ because e and f commute with K, see App. **A**.

We conclude that the lattice W-algebra $\mathcal{W}_{i,N}$ can be defined as an extension of the $\mathcal{TL}_{i,N}$ algebra by the generators

$$WB_j^+ = (-1)^j\big(c_j^\dagger c_{j+1}^\dagger + i c_j^\dagger c_{j+2}^\dagger - c_{j+1}^\dagger c_{j+2}^\dagger\big), \tag{13}$$

$$WB_j^0 = -\frac{1}{2}(1 + i c_j^\dagger c_{j+1} - c_j^\dagger c_{j+2} + i c_{j+1}^\dagger c_j - 2c_{j+1}^\dagger c_{j+1} - i c_{j+1}^\dagger c_{j+2} - c_{j+2}^\dagger c_j - i c_{j+2}^\dagger c_{j+1}), \tag{14}$$

$$WB_j^- = (-1)^{j+1}\big(c_jc_{j+1} + ic_jc_{j+2} - c_{j+1}c_{j+2}\big), \tag{15}$$

where $j = 1, 2, \ldots, N-2$. Of course, this statement should be proven. We formulate our result in Thm. **B.1** and give a computational proof in App. **B** where we use more convenient fermions (the Fourier transforms of c_j's and $c_j^\dagger$'s) introduced below.

We note that the relation of WB_j^0 generators with the wall Brauer generators (11) is

$$WB_j^0 = -\frac{1}{2}WB_j + \frac{i}{2}e_j - \frac{i}{2}e_{j+1}. \tag{16}$$

2.4.2. *$s\ell(2)$ structure.* The introduction of the WB_j^0 generators is convenient to have a triplet of generators with respect to the action of the $s\ell(2)$ part of $U_\mathfrak{q}s\ell(2)$ given in (9). Indeed, we note that the generators WB_j^+, WB_j^0 and WB_j^- for each site j form an $s\ell(2)$ triplet:

$$[\mathsf{e}, WB_j^+] = 0, \qquad [\mathsf{e}, WB_j^0] = -WB_j^+, \qquad [\mathsf{e}, WB_j^-] = 2WB_j^0, \tag{17}$$

$$[\mathsf{f}, WB_j^+] = -2WB_j^0, \qquad [\mathsf{f}, WB_j^0] = WB_j^-, \qquad [\mathsf{f}, WB_j^-] = 0, \tag{18}$$

and themselves satisfy $s\ell(2)$ relations on each site

$$[WB_j^+, WB_j^-] = 2WB_j^0, \qquad [WB_j^0, WB_j^\pm] = \pm WB_j^\pm.$$

We note that these statements agree with a more general statement given below in **4.7.1** for all roots of unity.

The formulas (17) and (18) mean that we have an action of $s\ell(2)$ on $\mathcal{W}_{i,N}$ by derivatives of the multiplication. Because the generators belong to spin-0 and spin-1 representations of the $s\ell(2)$, the algebra $\mathcal{W}_{i,N}$ is therefore decomposed onto $s\ell(2)$-modules of integer spins only. The multiplicity m_k of k-dimensional $s\ell(2)$ submodule in $\mathcal{W}_{i,N}$ is

$$m_k = \binom{2N-3}{N-k} + \binom{2N-3}{N-k-1} - \binom{2N-3}{N-k-2} - \binom{2N-3}{N-k-3} = \\ = \binom{2N-2}{N-k} - \binom{2N-2}{N-k-2}, \tag{19}$$

where k is odd. We note that m_1 is the Catalan number which equals the dimension of the TL subalgebra. This agrees with the fact that the span of $U_\mathfrak{q}s\ell(2)$-invariants is the subalgebra $\mathcal{TL}_{i,N}$. We finally obtain the dimension of the $\mathcal{W}_{i,N}$ algebra

$$\dim(\mathcal{W}_{i,N}) = \sum_{k\text{–odd}} km_k = 2^{2N-3}. \tag{20}$$

The numbers m_k and the dimension of the algebra can be computed explicitly using the representation theory of $\mathcal{W}_{i,N}$ which will be discussed briefly below. We just note now that the number 2^{2N-3} obtained as the dimension of the algebra follows from the fact that the dimension of its centralizer – $\rho_{g\ell}(\overline{U}_\mathfrak{q}s\ell(2))$ – is $8 = 2^3$ while the full matrix algebra (or the Clifford algebra) has dimension 2^{2N}.

2.4.3. *Relations in the algebra.* It would be interesting of course to express our algebra $\mathcal{W}_{i,N}$ directly in terms of generators and defining relations, much as can be done for the Temperley–Lieb algebra, but we have not managed to do so. Investigation of what the relations involving *WB* generators might be is quite intricate, and can be organized in terms of the number of sites involved. We give examples of the relations in App. **C**. We hope that there exist a bigger algebra with simpler defining relations (like for TL or Hecke algebras) such that its spin-chain representation is not faithful and this algebra factors through $\mathcal{W}_{i,N}$.

2.5. **Fermionic modes.**

2.5.1. *Hamiltonian spectrum and Jordan blocks.* In order to study the behavior of the lattice W-algebra in the continuum limit, we have to first discuss technical preliminaries. Fourier transform will play an essential role in what follows, and corresponding definitions are better understood by first studying here the spectrum, eigenvectors and Jordan blocks structure of the Hamiltonian in (6). We first compute eigenvectors of the adjoint action of H in the one-particle sector with the result

$$[H,\theta_k^\dagger]=2\cos\pi\frac{k}{N}\theta_k^\dagger,\qquad [H,\theta_k]=-2\cos\pi\frac{k}{N}\theta_k, \tag{21}$$

where we introduce fermions in the momentum space

$$\theta_k^\dagger=\sum_{j=1}^N A_k(j)c_j^\dagger,\qquad \theta_k=-i\sum_{j=1}^N A_k(j)c_j,\qquad k=1,\dots,N-1 \tag{22}$$

with the amplitudes

$$A_k(j)=\frac{1}{2}(1+ie^{-i\pi\frac{k}{N}})e^{i\pi\frac{kj}{N}}-\frac{1}{2}(1+ie^{i\pi\frac{k}{N}})e^{-i\pi\frac{kj}{N}},\qquad k=1,\dots,N-1. \tag{23}$$

These operators satisfy the anti-commutation relations

$$\{\theta_k^\dagger,\theta_{k'}\}=N\cos\pi\frac{k}{N}\delta_{k,k'}. \tag{24}$$

There are also two root vectors

$$\gamma^+=i\frac{\sqrt{\pi}}{N}\sum_{j=1}^N\Big(\frac{N}{2}-j+\frac{1}{2}\big(1-(-1)^j\big)\Big)i^jc_j^\dagger, \tag{25}$$

$$\gamma^-=-\frac{\sqrt{\pi}}{N}\sum_{j=1}^N\Big(\frac{N}{2}-j+\frac{1}{2}\big(1-(-1)^j\big)\Big)i^jc_j \tag{26}$$

which transform under the action of H as

$$[H,\gamma^+]=-\frac{2\sqrt{\pi}}{N}\theta_{N/2}^\dagger,\qquad [H,\gamma^-]=-\frac{2\sqrt{\pi}}{N}\theta_{N/2} \tag{27}$$

and satisfy the relations

$$\{\theta_{N/2}^\dagger,\gamma^-\}=\sqrt{\pi},\qquad \{\theta_{N/2},\gamma^+\}=\sqrt{\pi}. \tag{28}$$

Using the spectrum (21) of H, we conclude that the vacuum or ground state is

$$|0\rangle = \theta^\dagger_{N/2}\theta^\dagger_{N/2+1}\dots\theta^\dagger_{N-1}|\downarrow\dots\downarrow\rangle \tag{29}$$

and the log-partner of the vacuum is

$$\widetilde{|0\rangle} = \gamma^+\theta^\dagger_{N/2+1}\dots\theta^\dagger_{N-1}|\downarrow\dots\downarrow\rangle, \tag{30}$$

where $|\downarrow\dots\downarrow\rangle$ is a reference state with all spins down.

2.5.2. *The spectrum generating algebra.* For later convenience in the scaling limit section, we introduce new fermions

$$\begin{aligned}\eta^+_n &= \sqrt{\tfrac{n}{N\sin\pi\frac{n}{N}}}\theta^\dagger_{N/2+n},\\ \eta^-_n &= \sqrt{\tfrac{n}{N\sin\pi\frac{n}{N}}}\theta_{N/2-n},\end{aligned}\qquad n=-\frac{N}{2}+1,\dots,\frac{N}{2}-1,\quad n\neq 0, \tag{31}$$

together with the fermionic zero modes

$$\eta^+_0 = \frac{1}{\sqrt{\pi}}\theta^\dagger_{N/2},\qquad \eta^-_0 = \frac{1}{\sqrt{\pi}}\theta_{N/2}. \tag{32}$$

We have for these generators the anti-commutation relations

$$\{\eta^+_n,\eta^-_{n'}\} = n\delta_{n+n',0},\qquad \{\eta^\pm_0,\gamma^\mp\} = \pm 1. \tag{33}$$

The Hamiltonian (6) takes then the normal ordered form

$$H = \sum_{n=1}^{N/2-1}\frac{2}{n}\sin\pi\frac{n}{N}\big(\eta^-_{-n}\eta^+_n - \eta^+_{-n}\eta^-_n\big) - \frac{2\pi}{N}\eta^+_0\eta^-_0 + 1 - \cot\frac{\pi}{2N}, \tag{34}$$

with the groud state energy

$$H|0\rangle = (1-\cot\frac{\pi}{2N})|0\rangle, \tag{35}$$

where the vacuum now satisfies

$$\eta^\pm_{\geq 0}|0\rangle = 0. \tag{36}$$

We call the algebra generated by these fermions the spectrum generating algebra. Note that the adjoint action by the Hamiltonian is

$$[H,\eta^\pm_n] = -2\sin\pi\frac{n}{N}\eta^\pm_n,\quad [H,\gamma^\pm] = \frac{2\pi}{N}\eta^\pm_0. \tag{37}$$

2.5.3. *Quantum-group generators in terms of Fourier modes.* The quantum group generators introduced in (8) and (9) are expressed in terms of the $\eta^{\pm}$-fermions as

$$\mathsf{E} = \eta_0^+ \mathsf{K}, \qquad \mathsf{F} = \eta_0^- \tag{38}$$

with the $s\ell(2)$ generators

$$\mathsf{e} = \sum_{n=1}^{N/2-1} \frac{1}{n}\eta_{-n}^+\eta_n^+ + \gamma^+\eta_0^+, \qquad \mathsf{f} = -\sum_{n=1}^{N/2-1} \frac{1}{n}\eta_{-n}^-\eta_n^- - \gamma^-\eta_0^-, \tag{39}$$

$$\mathsf{h} = -\frac{1}{2}\Big(\sum_{n=1}^{N/2-1} \frac{1}{n}\big(\eta_{-n}^+\eta_n^- + \eta_{-n}^-\eta_n^+\big) + \gamma^-\eta_0^+ + \gamma^+\eta_0^-\Big). \tag{40}$$

We then have the $s\ell(2)$ action on the fermions

$$[\mathsf{e}, \eta_n^-] = \eta_n^+, \quad [\mathsf{f}, \eta_n^+] = \eta_n^-, \quad [\mathsf{h}, \eta_n^\pm] = \pm\frac{1}{2}\eta_n^\pm, \tag{41}$$

$$[\mathsf{e}, \gamma^-] = \gamma^+, \quad [\mathsf{f}, \gamma^+] = \gamma^-, \quad [\mathsf{h}, \gamma^\pm] = \pm\frac{1}{2}\gamma^\pm. \tag{42}$$

2.5.4. *TL and W generators in terms of Fourier modes.* Using the transformation (22), the TL generators (7) in terms of fermions (31) take the form

$$e_k = \frac{4(-1)^{n+1}}{N} \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin\pi j_1/N}{j_1}}\sqrt{\frac{\sin\pi j_2/N}{j_2}} \sin\pi k\Big(\frac{1}{2}+\frac{j_1}{N}\Big)\sin\pi k\Big(\frac{1}{2}+\frac{j_2}{N}\Big)\eta_{j_1}^+\eta_{j_2}^-, \tag{43}$$

where (and in what follows) we assume

$$\frac{\sin\pi j/N}{j}\Big|_{j=0} = \frac{\pi}{N}.$$

Meanwhile, the generators WB_j^α (13) of the algebra $\mathcal{W}_{i,N}$ can be written as

$$WB_k^+ = \frac{2}{N} \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin\pi j_1/N}{j_1}}\sqrt{\frac{\sin\pi j_2/N}{j_2}} V_k(j_1,j_2)\eta_{j_1}^+\eta_{j_2}^+, \tag{44}$$

$$WB_k^0 = \frac{2}{N} \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin\pi j_1/N}{j_1}}\sqrt{\frac{\sin\pi j_2/N}{j_2}} V_k(j_1,j_2)\big(\eta_{j_1}^+\eta_{j_2}^- + \eta_{j_1}^-\eta_{j_2}^+\big), \tag{45}$$

$$WB_k^- = \frac{2}{N} \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin\pi j_1/N}{j_1}}\sqrt{\frac{\sin\pi j_2/N}{j_2}} V_k(j_1,j_2)\eta_{j_1}^-\eta_{j_2}^-, \tag{46}$$

where we introduce trigonometric functions

$$V_k(j_1,j_2) = \sin\pi\frac{j_1-j_2}{2N}\cos\pi\frac{(j_1+j_2)(k+\frac{1}{2})}{N} - (-1)^k\cos\pi\frac{j_1+j_2}{2N}\sin\pi\frac{(j_1-j_2)(k+\frac{1}{2})}{N}.$$

We note that these functions have the symmetries

$$V_k(-j_1,-j_2) = V_k(j_2,j_1) = -V_k(j_1,j_2).$$

2.6. **Generalized lattice modes.** Our ultimate purpose is to establish a connection between the lattice algebras and the ones present in the corresponding conformal field theory. The intuition guiding us is that the Temperley–Lieb generators are in some sense a lattice version of the stress energy tensor $T(z)$, and similarly the WB_j^α generators a lattice version of the triplet chiral W-algebra currents $W^\alpha(z)$. The strategy to make this more precise is to define Fourier modes expected to converge in a certain sense (to be discussed below) towards the Virasoro L_m and W-algebra modes W_n^α. If such a convergence occurs, one might expect that the lattice modes obey, in the limit $N \to \infty$, the commutation relations of the CFT.

Of course, the algebra of lattice modes will not in general be closed, and calculating commutators will generate more quantities on the right hand side which should also converge to the Virasoro and W-algebra modes. This means that there must be a most likely infinite family of lattice versions of the L_m and W_n^α, with convergence to a single Virasoro and W-algebra occurring in the limit $N \to \infty$. In this subsection, we define this family explicitely, postponing the discussion of what happens as $N \to \infty$ to next section.

2.6.1. *Higher approximation of Virasoro modes.* The fact that many lattice expressions can converge to the same continuum limit is well known. For instance, there is an infinite number of choices of natural hamiltonians for the XXZ spin-chains coming out of the Quantum Inverse Scattering formalism which all go over to the Virasoro dilatation operator L_0 in the continuum limit. The underlying construction [10] suggests a natural extension to the case of non-zero modes – that is Fourier modes at non vanishing momentum.

We start by defining the operators

$$H_n^0 = \sum_{j=1}^{N-1} \cos \pi \frac{nj}{N} e_j, \qquad n \in \mathbb{Z}. \tag{47}$$

The Hamiltonian (6) in this notation is H_0^0. The adjoint action by the Hamiltonian H_0^0 on H_n^0 generates the family H_n^r with $r \in \mathbb{N}_0$ and $n \in \mathbb{Z}$ in the following way

$$[H_0^0, H_m^r] = -4 \sin \pi \frac{m}{2N} H_m^{r+1}, \qquad r = 0,1,2,\dots. \tag{48}$$

We note that this recurrence does not determine H_0^r directly but neatly means that H_0^0 commutes with H_0^r. To define H_0^r we calculate H_m^r with $m \neq 0$ using the recurrence and then set $m = 0$ in the final formula. Relation (47) bears some similarity with the definition of ladder operators in integrable XXZ spin chains [24], but is different.

We recall then the fermionic expression for the Hamiltonian in (34). By a direct calculation using the formulas (43), and (47), and the recurrence relation (48), we obtain

$$H_n^r = \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin \pi j_1/N}{j_1}}\sqrt{\frac{\sin \pi j_2/N}{j_2}}\times \tag{49}$$
$$\times \Big(\cos^r \pi\frac{j_1-j_2}{2N}\big((-1)^r\delta_{j_1+j_2,n}+\delta_{j_2+j_1,-n}\big)-$$
$$-\sin^r \pi\frac{j_1+j_2}{2N}\big(\delta_{j_1-j_2,N+n}+\delta_{j_1-j_2,-N-n}+(-1)^r\delta_{j_1-j_2,N-n}+(-1)^r\delta_{j_1-j_2,-N+n}\big)\Big)\eta^+_{j_1}\eta^-_{j_2}.$$

That these operators give in the scaling limit the Virasoro modes and more generally elements in the universal enveloping of a Virasoro algebra will be discussed below in the next section. Now, we introduce what will turn out later to be lattice analogues of modes of the W-algebra currents.

2.6.2. *Higher approximations of W-algebra modes.* Similarly, we first define the operators

$$W_n^{\alpha,0} = \sum_{j=1}^{N-2} \cos \pi\frac{n(j+\frac12)}{N} WB_j^\alpha, \qquad \alpha=+,-,0. \tag{50}$$

The adjoint action by the Hamiltonian H_0^0 on $W_n^{\alpha,0}$ generates the family $W_n^{\alpha,r}$ with $r\in\mathbb{N}_0$ and $n\in\mathbb{Z}$ in the following way

$$[H_0^0, W_m^{\alpha,r}] = -4\sin\pi\frac{m}{2N}W_m^{\alpha,r+1}, \qquad r=0,1,2,\dots, \qquad \alpha=+,-,0. \tag{51}$$

We note that $W_0^{\alpha,r}$ is determined in the same way as it is explained after (48).

Using the formula (49) for H_0^0, formulas (44)–(46) and the recurrent relation (51) we obtain

$$W_n^{+,r} = \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin \pi j_1/N}{j_1}}\sqrt{\frac{\sin \pi j_2/N}{j_2}} U_{r,n}(j_1,j_2)\eta^+_{j_1}\eta^+_{j_2}, \tag{52}$$

$$W_n^{0,r} = \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin \pi j_1/N}{j_1}}\sqrt{\frac{\sin \pi j_2/N}{j_2}} U_{r,n}(j_1,j_2)\Big(\eta^+_{j_1}\eta^-_{j_2}+\eta^-_{j_1}\eta^+_{j_2}\Big), \tag{53}$$

$$W_n^{-,r} = \sum_{j_1,j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \sqrt{\frac{\sin \pi j_1/N}{j_1}}\sqrt{\frac{\sin \pi j_2/N}{j_2}} U_{r,n}(j_1,j_2)\eta^-_{j_1}\eta^-_{j_2}, \tag{54}$$

where we introduce trigonometric functions

$$U_{r,n}(j_1,j_2) = \sin\pi\frac{j_1-j_2}{2N}\cos^r\pi\frac{j_1-j_2}{2N}\big((-1)^r\delta_{j_1+j_2,n}+\delta_{j_1+j_2,-n}\big) \tag{55}$$
$$+2\cos\pi\frac{j_1+j_2}{2N}\sin^r\pi\frac{j_1+j_2}{2N}\big(\delta_{j_1-j_2,-N-n}+(-1)^r\delta_{j_1-j_2,-N+n}\big).$$

2.7. **A note on the odd number of sites.** We can also consider the spin-chains with odd number N of sites. They will give below a different sector of symplectic fermions in the scaling limit.

For the odd-N we define the fermions $\eta_n^\pm$ with half-integer n by formulas (31) where fermions θ_k and $\theta_k^\dagger$ are defined by (22) with amplitudes $A_k(j)$ defined by (23), for $k = 1, \dots, N-1$, and $A_0(j) = i^j$. We also have the generators (8) of the quantum group, which satisfy

$$\{\mathsf{EK}^{-1}, \eta_j^\pm\} = 0, \quad \{\mathsf{F}, \eta_j^\pm\} = 0, \quad j = -N/2+1, -N/2+2, \dots, N/2-1 \tag{56}$$

in addition to the quantum group relations. Then the backward transformation is

$$c_k^\dagger = -i \sum_{j=-N/2+1}^{N/2-1} \frac{1}{j} \sqrt{\frac{j}{N \sin \pi j/N}} A_{N/2+j}(k) \eta_j^+ - i^k \mathsf{EK}^{-1}, \tag{57}$$

$$c_k = - \sum_{j=-N/2+1}^{N/2-1} \frac{1}{j} \sqrt{\frac{j}{N \sin \pi j/N}} A_{N/2-j}(k) \eta_j^- + i^{k-1} \mathsf{F}. \tag{58}$$

The $s\ell(2)$ generators are now represented as

$$\mathsf{e} = \sum_{n=1/2}^{N/2-1} \frac{1}{n} \eta_{-n}^+ \eta_n^+, \qquad \mathsf{f} = - \sum_{n=1/2}^{N/2-1} \frac{1}{n} \eta_{-n}^- \eta_n^-,$$

$$\mathsf{h} = -\frac{1}{2} \sum_{n=1/2}^{N/2-1} \frac{1}{n} \big(\eta_{-n}^+ \eta_n^- + \eta_{-n}^- \eta_n^+ \big).$$

We note that the Hamiltonian (6) takes a slightly different form

$$H = \sum_{n=1/2}^{N/2-1} \frac{2}{n} \sin \pi \frac{n}{N} \big(\eta_{-n}^- \eta_n^+ - \eta_{-n}^+ \eta_n^- \big) + \left(1 - \frac{1}{\sin \frac{\pi}{2N}} \right). \tag{59}$$

The definition of the lattice W-algebra is the same as in the even-N case and Thm. **B.1** is true for odd-N case as well. Formal expressions of the lattice W-algebra generators in terms of $\eta^\pm$ fermions are the same as in the even-N case. So, the generators e_k are given by (43) and W_k^α are given by (44)-(46). Similarly, their linear combinations H_n^0 and $W_n^{\alpha,0}$ together higher "currents" H_n^r and $W_n^{\alpha,r}$ are expressed in the same way. So, the currents H_n^r are given by (49) and currents $W_n^{\alpha,r}$ by (52)–(54). For odd N the operators H_n^r and $W_n^{\alpha,r}$ satisfy the same commutation relations as for even N.

Next, we turn to the study of the scaling properties of the whole family of the operators (modes) H_n^r and $W_n^{\alpha,r}$ introduced in **2.6**.

3. Scaling limit of the XX chain and W-algebra modes

In this section we calculate the limit $N \to \infty$ of the commutation relations between currents (49) and (52)–(54) and show that they coincide with the commutation relations of the triplet W-algebra. For the convenience of the reader we give all needed conventions about symplectic fermions and the triplet W-algebra in App. **D**.

3.1. Scaling limit and $1/N$-decomposition.

Here, we discuss how to proceed from the $\mathcal{W}_{i,N}$ generators to get all the modes of the W-algebra currents, including the Virasoro generators, in the chiral logarithmic conformal field theory of symplectic fermions. The Fourier modes H_n^r and $W_n^{\alpha,r}$ introduced in the previous section are expressed as fermionic bilinears with coefficients being functions of the spin-chain size N. We will show below that they generate a finite dimensional algebra whose structure constants are trigonometric functions of N. The scaling limit of the Fourier modes and their commutators involves formal expansions of the operators and the structure constants, respectively, as Laurent series in N. Differnt terms in such expansions for H_n^r and $W_n^{\alpha,r}$ are then identified with elements from the enveloping algebra of the chiral triplet W-algebra.

3.1.1. *The scaling limit of the Hamiltonian.* We first study the scaling limit of the Hamiltonian H. We take the Hamiltonian (34), which is written in the normal-ordered form and keep terms up to the order $1/N$ in the $N \to \infty$ decomposition

$$H = -2\frac{N}{\pi} + 1 + 2\frac{\pi}{N}\Big(-\frac{(-2)}{24} + \sum_{n=1}^{N/2-1}\big(\eta_{-n}^-\eta_n^+ - \eta_{-n}^+\eta_n^-\big) - \eta_0^+\eta_0^-\Big) + O(1/N^2). \tag{60}$$

We see that the leading part (after subtracting the vacuum expectation of the Hamiltonian) which is an operator in front of $1/N$ has the form $L_0 - c/24$ on the scaling states, with $c = -2$. This expression of L_0 is well-known and it is the zero mode of the stress-energy tensor $T(z)$ in the symplectic fermions theory.

3.1.2. *Higher hamiltonians.* We could expect new features considering the scaling limit of the whole family of the higher hamiltonians of the open XX spin-chain introduced above.

The limit $N \to \infty$ of operators H_0^{2k} given by (49) is singular and therefore we should subtract the divergent part. To do this we calculate the eigenvalues $2h_\ell$ of H_0^ℓ on the vacuum vector (29) using (49), which gives

$$h_0 = \frac{1}{2}\big(\cot\pi\frac{1}{2N} - 1\big), \tag{61}$$

$$h_\ell = -\frac{1}{2(\ell+1)} + \frac{1}{2^{\ell+1}}\sum_{n=0}^{\ell/2}\left(\binom{\ell-1}{\ell/2+n-1} - \binom{\ell-1}{\ell/2+n+1}\right)\cot\pi\frac{2n+1}{2N}, \quad \text{for even } \ell > 0.$$

3.1.3. *Virasoro modes in symplectic fermions.* We start from calculating the scaling limit of H_n^0. Decomposing $\sqrt{\sin \pi j/N}$ in (49) into a series in $\frac{1}{N}$ we obtain

$$\frac{N}{\pi}(H_n^0 - 2h_0\delta_{n,0}) = -\sum_{j=-\frac{N}{2}+1}^{\frac{N}{2}-1} :\eta_j^+\eta_{n-j}^-: - \sum_{j=-\frac{N}{2}+1}^{\frac{N}{2}-1} :\eta_j^+\eta_{-n-j}^-: +$$
$$+ \sum_{j_1=-\frac{N}{2}+1}^{\frac{N}{2}-1}\sum_{j_2=-\frac{N}{2}+1}^{\frac{N}{2}-1} \big(\delta_{j_1-j_2,N+n} + \delta_{j_1-j_2,-N-n} + \delta_{j_1-j_2,N-n} + \delta_{j_1-j_2,-N+n}\big)\eta_{j_1}^+\eta_{j_2}^- + o\big(\tfrac{1}{N}\big).$$

We note that the second line of the previous expression vanishes on the scaling states because for any n we can find N such that j_1 or j_2 will be larger than any fixed number. Therefore on scaling states we have

$$\frac{N}{\pi}(H_n^0 - 2h_0\delta_{n,0}) = L_n^{(N)} + L_{-n}^{(N)} + o\big(\tfrac{1}{N}\big), \tag{62}$$

where

$$L_n^{(N)} = -\sum_{j=-\frac{N}{2}+1}^{\frac{N}{2}-1} :\eta_j^+\eta_{n-j}^-: . \tag{63}$$

This expression coinsides with (183) in the limit $N \to \infty$. In what follows we suppress the superscript $^{(N)}$ in the expressions whenever it cannot lead to a confusion.

After subtracting the vacuum eigenvalues we can write the series decomposition of (49) in $\frac{1}{N}$ on scaling states

$$H_n^r - (1+(-1)^r)h_r\delta_{n,0} = \sum_{k=1}^{\infty}\Big(\frac{\pi}{N}\Big)^{2k-1} H_n^r[k], \tag{64}$$

where first several terms have the form

$$H_n^r[1] = (-1)^r L_n + L_{-n}, \tag{65}$$

$$H_n^r[2] = \frac{1}{24}\Big(4(3r+1)\big((-1)^r(L^2)_n + (L^2)_{-n}\big) - 3r\big((-1)^r(\partial^2 L)_n + (\partial^2 L)_{-n}\big) + \tag{66}$$
$$+ 3(r+2)\big((-1)^r(\partial L)_n + (\partial L)_{-n}\big) + 4\big((-1)^r L_n + L_{-n}\big)\Big),$$

$$H_n^r[3] = \frac{1}{5760}\Big(-64(15r^2+1)\big((-1)^r(L^3)_n + (L^3)_{-n}\big) + \tag{67}$$
$$+ 8(2+15r(r-1))\big((-1)^r(\partial^2 L^2)_n + (\partial^2 L^2)_{-n}\big) +$$
$$+ (-4+15r(r+2))\big((-1)^r(\partial^4 L)_n + (\partial^4 L)_{-n}\big) -$$
$$- 1080r(r+1)\big((-1)^r(\partial^2 L^2)_n + (\partial^2 L^2)_{-n}\big) +$$
$$+ 10(-4+3r(9r+8))\big((-1)^r(\partial^3 L)_n + (\partial^3 L)_{-n}\big) -$$
$$- 80(9r^2+24r-1)\big((-1)^r(L^2)_n + (L^2)_{-n}\big) +$$
$$+ 45(r^2+6r-4)\big((-1)^r(\partial^2 L)_n + (\partial^2 L)_{-n}\big) -$$

$$- 45(r^2+6r+4)\big((-1)^r(\partial L)_n + (\partial L)_{-n}\big) - \\ - 48\big((-1)^r L_n + L_{-n}\big)\Big),$$

where expressions for L_n and $(L^2)_n$ are given in (183) and (184).

In order to obtain the defining relations of the Virasoro algebra in the continuum limit, only the leading $1/N$ term is necessary, that is the $H_n^r[1]$. For the W algebra relations, the presence of non linearities requires keeping also $H_n^r[2]$, and similarly (see below) $WB_n^{\alpha,r}[1]$ and $WB_n^{\alpha,r}[2]$.

3.1.4. *W-algebra modes in symplectic fermions.* In a similar way we obtain the decompositions of (52)–(54)

$$W_n^{\alpha,r} = \sum_{k=1}^{\infty} \Big(\frac{\pi}{N}\Big)^{2k} W_n^{\alpha,r}[k], \tag{68}$$

where the first two coefficients are

$$W_n^{\alpha,r}[1] = (-1)^r W_n^\alpha + W_{-n}^\alpha, \tag{69}$$

$$W_n^{\alpha,r}[2] = \frac{1}{48}\Big(\frac{48}{5}(3r+2)\big((-1)^r(LW^\alpha)_n + (LW^\alpha)_{-n}\big) - \\ - \frac{2}{5}(9r+1)\big((-1)^r(\partial^2 W^\alpha)_n + (\partial^2 W^\alpha)_{-n}\big) - \\ - (18r+26)\big((-1)^r(\partial W^\alpha)_n + (\partial W^\alpha)_{-n}\big) - \\ - (6r+22)\big((-1)^r W_n^\alpha + W_{-n}^\alpha\big)\Big). \tag{70}$$

3.2. **Scaling limit of the commutators.** In this section we calculate commutators of H_n^r and $W_n^{\alpha,r}$ and show that in the limit $N\to\infty$ the commutators give the commutation relations (189)–(191) of the W-algebra. Using the explicit expression (49) we obtain the following commutation relations

$$[H_n^0, H_m^0] = 2\sin\pi\frac{n-m}{2N} H_{n+m}^1 + 2\sin\pi\frac{n+m}{2N} H_{n-m}^1, \tag{71}$$

$$[H_n^0, H_m^1] = 2\sin\pi\frac{2n-m}{2N} H_{n+m}^2 - 2\sin\pi\frac{2n+m}{2N} H_{n-m}^2 - \\ - 2\sin\pi\frac{n}{2N}\big(\cos\pi\frac{n-m}{2N} H_{n+m}^0 - \cos\pi\frac{n+m}{2N} H_{n-m}^0\big), \tag{72}$$

$$[H_n^1, H_m^1] = 2\sin\pi\frac{n-m}{N} H_{n+m}^3 - 2\sin\pi\frac{n+m}{N} H_{n-m}^3 - \\ - 2\sin\pi\frac{n-m}{2N}(\cos\pi\frac{n+m}{2N} + \sin\pi\frac{n}{2N}\sin\pi\frac{m}{2N}) H_{n+m}^1 + \\ + 2\sin\pi\frac{n+m}{2N}(\cos\pi\frac{n-m}{2N} - \sin\pi\frac{n}{2N}\sin\pi\frac{m}{2N}) H_{n-m}^1. \tag{73}$$

Taking (64) and (65) into account we obtain the commutation relations of L_n as the limit

$$[L_n, L_m] = \lim_{N\to\infty} \frac{N^2}{4\pi^2}\big([H^0_n, H^0_m] - [H^0_n, H^1_m] + [H^0_m, H^1_n] + [H^1_n, H^1_m]\big). \tag{74}$$

Then in the previous formula we substitute right hand sides of (71)–(73), then substitute H^r_n from (64) and calculate the Laurent series in N up to the second order which gives the Virasoro commutation relations (189). We note that in this calculation we need to keep only first term in the decomposition (64). We also note that the Virasoro central extension appears from the decomposition of the vacuum eigenvalues (61) in N.

The commutation relations $[H^r_n, W^{\alpha,s}_m]$ and $[W^{\alpha,r}_n, W^{\beta,s}_m]$ are given in App. **E**. To calculate the commutators

$$[L_n, W^\alpha_m] = \lim_{N\to\infty} \frac{N^3}{4\pi^3}\big([H^0_n, W^{\alpha,0}_m] - [H^0_n, W^{\alpha,1}_m] - [H^1_m, W^{\alpha,0}_m] + [H^1_n, W^{\alpha,1}_m]\big) \tag{75}$$

we take the commutators from (192)–(195) and substitute the Laurent decompositions (64) and (68) into them. This gives (190).

The calculation of the commutator

$$[W^\alpha_n, W^\beta_m] = \lim_{N\to\infty} \frac{N^4}{4\pi^4}\big([W^{\alpha,0}_n, W^{\beta,0}_m] - [W^{\alpha,1}_n, W^{\beta,1}_m] - [W^{\alpha,1}_m, W^{\beta,0}_m] + [W^{\alpha,1}_n, W^{\beta,1}_m]\big) \tag{76}$$

differs from the previous two in the following point. Because we have the factor N^4 in front of the commutators we should take the decompositions of the right hand sides of (196)–(198) up to the 4th order. The leading order coming from the trigonometric functions is 0 or 1 and therefore we should keep the 3rd and 4th orders in the decompositions (64) and (68) respectively. This calculation results in the contribution from $(L^2)_n$ and $(LW^\alpha)_n$ in (191), which is obtained after taking the limit. We have thus succeeded in reproducing the algebra of Virasoro and W modes from our lattice approximations.

3.3. **A note on the odd-N case scaling limit.** The calculation of the $N \to \infty$ limit for the odd N differs from the calculation for the even N only by normal ordering constants. Thus for the odd case the calculation reapeats the even case where we take normal ordering constants

$$h_0 = \frac{1}{2}\left(\frac{1}{\sin\pi\frac{1}{2N}} - 1\right), \tag{77}$$

$$h_\ell = -\frac{1}{2(\ell+1)} + \frac{1}{2^{\ell+1}}\sum_{n=0}^{\ell/2}\left(\binom{\ell-1}{\ell/2+n-1} - \binom{\ell-1}{\ell/2+n+1}\right)\frac{1}{\sin\pi\frac{2n+1}{2N}}, \quad \text{for even } \ell > 0$$

instead of (61).

4. The XXZ representation and the W-algebra on a lattice

In this section, we give our definition of the lattice W-algebra in the context of XXZ spin-chains at any root of unity $\mathfrak{q}$. We begin with collecting basic facts about XXZ spin-chains with quantum-group symmetry [23, 3, 32]. Then, we define the lattice W-algebra using its centralizing property with the small quantum group and study its properties (generators, relations, representation theory, etc.) Finally, in the last subsection we discuss two scaling limits of this lattice algebra, describing $(p-1,p)$ and $(1,p)$ models.

4.1. **The Hamiltonian.** We consider the XXZ Hamiltonian of N one-half spins with an open boundary condition described by the "quantum-group symmetric" boundary term [23],

$$H(\mathfrak{q}) = \frac{1}{2}\sum_{i=1}^{N-1}(\sigma_i^x\sigma_{i+1}^x + \sigma_i^y\sigma_{i+1}^y + \frac{\mathfrak{q}+\mathfrak{q}^{-1}}{2}\sigma_i^z\sigma_{i+1}^z) - \frac{\mathfrak{q}-\mathfrak{q}^{-1}}{4}(\sigma_1^z - \sigma_N^z), \tag{78}$$

where σ_i^x, σ_i^y and σ_i^z are usual Pauli matrices given in (3). We note that we have an opposite sign in front of the boundary term (78), with respect to a usual choice [23, 3], which corresponds to a slightly different basis (corresponding to $\mathfrak{q} \to \mathfrak{q}^{-1}$ or relabeling of sites from N to 1) we use in order to adapt to our quantum-group notations.

The Hamiltonian $H(\mathfrak{q})$ is not hermitian due to the imaginary boundary term $\frac{\sin(\pi/p)}{2\mathrm{i}}$ but its eigenvalues are real while its Jordan form has cells of rank 2. An easiest way to see non-trivial Jordan forms in the spectrum is to study symmetries of the Hamiltonian densities which we discuss below.

We note finally that the hamiltonian is massless when $|\mathfrak{q}| = 1$ and we will consider only this critical case. Moreover, roots of unity cases are most interesting for applications in the scaling limit and these are in our interest for definition and study lattice versions of the triplet W-algebras as well. We first remind basic information about the quantum group $U_{\mathfrak{q}}s\ell(2)$ symmetry at the root of unity cases.

4.2. **Quantum groups.** Here, we collect definitions of different quantum groups at roots of unity and recall basic facts about their representation theory. In setting the notation and recalling the basic facts about $U_{\mathfrak{q}}s\ell(2)$ needed below, we largely follow [25].

4.2.1. *The small quantum group.* The quantum group $\overline{U}_{\mathfrak{q}}s\ell(2)$ is the "small" quantum $s\ell(2)$ with $\mathfrak{q} = e^{i\pi/p}$ and the three generators E, F, and $\mathsf{K}^{\pm1}$ satisfying the standard relations for the quantum $s\ell(2)$,

$$\mathsf{K}\mathsf{E}\mathsf{K}^{-1} = \mathfrak{q}^2\mathsf{E}, \quad \mathsf{K}\mathsf{F}\mathsf{K}^{-1} = \mathfrak{q}^{-2}\mathsf{F}, \quad [\mathsf{E},\mathsf{F}] = \frac{\mathsf{K}-\mathsf{K}^{-1}}{\mathfrak{q}-\mathfrak{q}^{-1}}, \tag{79}$$

with some additional constraints,

$$\mathsf{E}^p = \mathsf{F}^p = 0, \quad \mathsf{K}^{2p} = \mathbf{1}, \tag{80}$$

and the comultiplication is given by

$$\Delta(\mathsf{E}) = \mathbf{1}\otimes\mathsf{E} + \mathsf{E}\otimes\mathsf{K}, \quad \Delta(\mathsf{F}) = \mathsf{K}^{-1}\otimes\mathsf{F} + \mathsf{F}\otimes\mathbf{1}, \quad \Delta(\mathsf{K}) = \mathsf{K}\otimes\mathsf{K}. \tag{81}$$

This associative algebra is finite-dimensional

4.2.2. *The full quantum group.* The Lusztig (or full) quantum group $U_{\mathfrak{q}}s\ell(2)$ with $\mathfrak{q} = e^{i\pi/p}$, for any integer $p \geq 2$, is generated by E, F, and $\mathsf{K}^{\pm 1}$ satisfying the relations (79) and (80), and additionally by the divided powers $\mathsf{f} \sim \mathsf{F}^p/[p]!$ and $\mathsf{e} \sim \mathsf{E}^p/[p]!$ which satisfy the usual $s\ell(2)$-relations:

$$[\mathsf{h},\mathsf{e}] = \mathsf{e}, \qquad [\mathsf{h},\mathsf{f}] = -\mathsf{f}, \qquad [\mathsf{e},\mathsf{f}] = 2\mathsf{h}.$$

The full list of relations with comultiplication formulae are listed in App. **A** where we also give relations to more usual (in the spin-chain literature) quantum group generators $S^{\pm}$, S^z and $\mathfrak{q}^{S^z}$.

4.3. **The Casimir operator and Temperley–Lieb algebra.** As a module over $U_{\mathfrak{q}}s\ell(2)$, the XXZ spin chain $\mathcal{H}_N$ is a tensor product of N copies of the "fundamental" two-dimensional simple module $\mathsf{X}_{2,1}$ (see notations below in **4.4.1**) such that the genereators are represented on $\mathsf{X}_{2,1}$ as

$$\mathsf{E} = \sigma^+, \quad \mathsf{F} = \sigma^-, \quad \mathsf{K} = \frac{\mathfrak{q}+\mathfrak{q}^{-1}}{2}\mathbf{1} + \frac{\mathfrak{q}-\mathfrak{q}^{-1}}{2}\sigma^z, \quad \text{and} \quad \mathsf{e} = \mathsf{f} = \mathsf{h} = 0.$$

Using the $(N-1)$-folded comultiplications (145) for the capital generators together with the general comultiplication (146) and (147) for the divided powers, which we compute in App. **A**, we obtain the XXZ representation of $U_{\mathfrak{q}}s\ell(2)$

$$\rho_{\mathfrak{q},N} : U_{\mathfrak{q}}s\ell(2) \to \mathrm{End}_{\mathbb{C}}(\mathcal{H}_N), \qquad \mathcal{H}_N = \bigotimes_{j=1}^{N} \mathsf{X}_{2,1}. \tag{82}$$

It has the usual XXZ expressions for the generators

$$\begin{aligned} \rho_{\mathfrak{q},N}(\mathsf{E}) &= \sum_{j=1}^{N} \mathbf{1}\otimes\ldots\otimes\mathbf{1}\otimes\sigma_j^+\otimes\mathfrak{q}^{\sigma_{j+1}^z}\otimes\ldots\otimes\mathfrak{q}^{\sigma_N^z}, \\ \rho_{\mathfrak{q},N}(\mathsf{F}) &= \sum_{j=1}^{N} \mathfrak{q}^{-\sigma_1^z}\otimes\ldots\otimes\mathfrak{q}^{-\sigma_{j-1}^z}\otimes\sigma_j^-\otimes\mathbf{1}\otimes\ldots\otimes\mathbf{1} \end{aligned} \tag{83}$$

and for the divided powers the action is

$$\begin{aligned} \rho_{\mathfrak{q},N}(\mathsf{e}) = \mathfrak{q}^{\frac{p(p-1)}{2}}\Big(\bigotimes_{j=1}^{N}\sigma_j^z\Big) \sum_{1\leq j_1<j_2<\cdots<j_p\leq N} (-1)^{j_p}\mathbf{1}\otimes\ldots\otimes\mathbf{1}\otimes\sigma_{j_1}^+\otimes\mathfrak{q}^{\sigma_{j_1+1}^z}\otimes\mathfrak{q}^{\sigma_{j_1+2}^z}\otimes\ldots \\ \ldots\otimes\mathfrak{q}^{(k-1)\sigma_{j_k-1}^z}\otimes\sigma_{j_k}^+\mathfrak{q}^{(k-1)\sigma_{j_k}^z}\otimes\mathfrak{q}^{k\sigma_{j_k+1}^z}\otimes\ldots\otimes\sigma_{j_p}^+\mathfrak{q}^{(p-1)\sigma_{j_p}^z}\otimes\sigma_{j_p+1}^z\otimes\ldots\otimes\sigma_N^z \end{aligned} \tag{84}$$

and

$$\rho_{\mathfrak{q},N}(\mathsf{f}) = (-1)^p\Big(\bigotimes_{j=1}^{N}\sigma_j^z\Big)\sum_{1\le j_1<j_2<\dots<j_p\le N}(-1)^{j_p}\mathbf{1}\otimes\dots\otimes\mathbf{1}\otimes\sigma_{j_1}^-\otimes\mathfrak{q}^{\sigma_{j_1+1}^z}\otimes\mathfrak{q}^{\sigma_{j_1+2}^z}\otimes\dots \\ \dots\otimes\mathfrak{q}^{(k-1)\sigma_{j_k-1}^z}\otimes\sigma_{j_k}^-\otimes\mathfrak{q}^{k\sigma_{j_k+1}^z}\otimes\dots\otimes\sigma_{j_p}^-\otimes\sigma_{j_p+1}^z\otimes\dots\otimes\sigma_N^z. \tag{85}$$

Then, we recall that the Casimir element

$$\boldsymbol{C} = \mathsf{F}\mathsf{E} + \frac{\mathfrak{q}\mathsf{K}+\mathfrak{q}^{-1}\mathsf{K}^{-1}}{(\mathfrak{q}-\mathfrak{q}^{-1})^2} \tag{86}$$

commutes with the full quantum group $U_\mathfrak{q}s\ell(2)$ and it is represented on $\mathsf{X}_{2,1}\otimes\mathsf{X}_{2,1}$ as

$$\Delta(\boldsymbol{C}) = \frac{1}{2}\big(\sigma^x\otimes\sigma^x+\sigma^y\otimes\sigma^y+\frac{\mathfrak{q}+\mathfrak{q}^{-1}}{2}\sigma^z\otimes\sigma^z\big) \\ +\frac{\mathfrak{q}-\mathfrak{q}^{-1}}{4}\big(\mathbf{1}\otimes\sigma^z-\sigma^z\otimes\mathbf{1}\big)+\frac{3\mathfrak{q}^3+\mathfrak{q}+\mathfrak{q}^{-1}+3\mathfrak{q}^{-3}}{4(\mathfrak{q}-\mathfrak{q}^{-1})^2}\mathbf{1}\otimes\mathbf{1}.$$

This gives the usual XXZ coupling and the Hamiltonian (78) is expressed as a sum over the densities $\Delta(\boldsymbol{C})_{i,i+1}$ acting on ith and $(i+1)$th tensorands:

$$H(\mathfrak{q}) = \sum_{i=1}^{N-1}\Delta(\boldsymbol{C})_{i,i+1} - (N-1)\frac{3\mathfrak{q}^3+\mathfrak{q}+\mathfrak{q}^{-1}+3\mathfrak{q}^{-3}}{4(\mathfrak{q}-\mathfrak{q}^{-1})^2}.$$

Introducing the more convenient notation

$$e_i = -\Delta(\boldsymbol{C})_{i,i+1}+\frac{\mathfrak{q}^3+\mathfrak{q}^{-3}}{(\mathfrak{q}-\mathfrak{q}^{-1})^2}\mathbf{1}, \qquad 1\le i\le N-1, \tag{87}$$

the Hamiltonian takes the usual form [23, 3]

$$H(\mathfrak{q}) = -\sum_{i=1}^{N-1}e_i + (N-1)\frac{\mathfrak{q}+\mathfrak{q}^{-1}}{4}. \tag{88}$$

It is straightforward to check that the operators e_i, for $1\le i\le N-1$, satisfy the defining relations of the Temperley–Lieb algebra $\mathcal{TL}_{\mathfrak{q},N}$

$$e_i^2 = (\mathfrak{q}+\mathfrak{q}^{-1})e_i, \\ e_ie_{i\pm1}e_i = e_i, \\ e_ie_j = e_je_i, \quad |i-j|>1.$$

The relation (87) of the Temperley–Lieb generators e_i with the quantum-group Casimir operator and the coassociativity of the quantum-group comultiplication give the well-known result [23]

$$\big[\rho_{\mathfrak{q},N}\big(U_\mathfrak{q}s\ell(2)\big),\mathcal{TL}_{\mathfrak{q},N}\big] = 0. \tag{89}$$

Moreover, it was shown first in [21] that $\mathcal{TL}_{\mathfrak{q},N}$ is the centralizer of the representation of $U_\mathfrak{q}s\ell(2)$ on $\mathcal{H}_N$ and vice versa, i.e. they are mutual centralizers,

$$\mathcal{TL}_{\mathfrak{q},N}\cong\mathrm{End}_{U_\mathfrak{q}s\ell(2)}(\mathcal{H}_N), \qquad \rho_{\mathfrak{q},N}\big(U_\mathfrak{q}s\ell(2)\big)\cong\mathrm{End}_{\mathcal{TL}_{\mathfrak{q},N}}(\mathcal{H}_N), \tag{90}$$

for any $\mathfrak{q}$ at the root of unity cases.

It was also shown [21, 3] that the representation (87) is faithful and projective $\mathcal{TL}_{\mathfrak{q},N}$-modules are direct summands in a decomposition of the spin-chain, in particular. It is thus important to describe representation theory of its centralizer $U_{\mathfrak{q}}s\ell(2)$ for any root of unity first before going into details on the spin-chain decomposition.

4.4. **Modules over the quantum groups.** In this section, we describe few important $U_{\mathfrak{q}}s\ell(2)$- and $\overline{U}_{\mathfrak{q}}s\ell(2)$-modules which appear in decompositions of the spin-chains. We begin with reminding results on the simple modules and then describe their projective covers [25].

4.4.1. *Simple $U_{\mathfrak{q}}s\ell(2)$-modules.* A simple $U_{\mathfrak{q}}s\ell(2)$-module $\mathsf{X}_{s,r}$ is labeled by the pair (s,r), with $1 \leq s \leq p$ and $r \in \mathbb{N}$, and has the highest weights $(-1)^{r-1}\mathfrak{q}^{s-1}$ and $\frac{r-1}{2}$ with respect to K and h generators, respectively. The sr-dimensional module $\mathsf{X}_{s,r}$ is spanned by elements $\mathsf{a}_{n,m}$, $0 \leq n \leq s-1$, $0 \leq m \leq r-1$, where $\mathsf{a}_{0,0}$ is the highest-weight vector and the left action of the algebra on $\mathsf{X}_{s,r}$ is given by

$$\mathsf{K}\mathsf{a}_{n,m} = (-1)^{r-1}\mathfrak{q}^{s-1-2n}\mathsf{a}_{n,m}, \qquad \mathsf{h}\,\mathsf{a}_{n,m} = \frac{1}{2}(r-1-2m)\mathsf{a}_{n,m}, \tag{91}$$


$$\mathsf{E}\mathsf{a}_{n,m} = (-1)^{r-1}[n][s-n]\mathsf{a}_{n-1,m}, \qquad \mathsf{e}\,\mathsf{a}_{n,m} = m(r-m)\mathsf{a}_{n,m-1}, \tag{92}$$

$$\mathsf{F}\mathsf{a}_{n,m} = \mathsf{a}_{n+1,m}, \qquad \mathsf{f}\,\mathsf{a}_{n,m} = \mathsf{a}_{n,m+1}, \tag{93}$$

where we set $\mathsf{a}_{-1,m} = \mathsf{a}_{n,-1} = \mathsf{a}_{s,m} = \mathsf{a}_{n,r} = 0$. We note that the $\mathsf{X}_{s,r}$ restricted to the subalgebra $\overline{U}_{\mathfrak{q}}s\ell(2)$ is isomorphic to the r-fold direct sum of the simple $\overline{U}_{\mathfrak{q}}s\ell(2)$-module X_s^α with $\alpha = (-1)^{r-1}$ and the action of $\overline{U}_{\mathfrak{q}}s\ell(2)$-generators is given by the first column of (91)-(93).

4.4.2. *Projective covers.* The simple modules $\mathsf{X}_{s,r}$ have projective covers $\mathsf{P}_{s,r}$ with the subquotient structure

(94)



where $r \geq 2$. We also note that the $\mathsf{P}_{s,r}$ as a $\overline{U}_{\mathfrak{q}}s\ell(2)$-module is isomorphic to the r-fold direct sum of P_s^α with $\alpha = (-1)^{r-1}$. The action of the generators E, F, and K in P_s^α can be found in [26].

4.5. **XXZ decompositions.** We are now ready to show a decomposition of $\mathcal{H}_N = \otimes^N \mathsf{X}_{2,1}$ over $U_{\mathfrak{q}}s\ell(2)$. This decomposition over the quantum group for any root of unity and any N was explicitly written in [32] and we borrowed this result:

$$\mathcal{H}_N|_{U_\mathfrak{q}s\ell(2)} \cong \bigoplus_{\substack{s=N \bmod 2+1,\\ s+N=1 \bmod 2}}^{p-1} \dim\big(\mathcal{X}_{\frac{s-1}{2}}\big)\mathsf{X}_{s,1} \oplus \bigoplus_{r=1}^{r_m-1} \bigoplus_{\substack{s=0,\\ rp+s+N=1 \bmod 2}}^{p-1} \dim\big(\mathcal{X}_{\frac{rp+s-1}{2}}\big)\mathsf{P}_{p-s,r} \oplus \bigoplus_{\substack{s=0,\\ s+s_m=1 \bmod 2}}^{s_m+1} \dim\big(\mathcal{X}_{\frac{r_m p+s-1}{2}}\big)\mathsf{P}_{p-s,r_m}\,, \tag{95}$$

where we defined $N = r_m p + s_m$, for $r_m \in \mathbb{N}$ and $-1 \le s_m \le p-2$, and a multiplicity of each $\mathsf{P}_{p-s,r}$ and $\mathsf{X}_{s,1}$ equals $\mathsf{d}^0_{\frac{rp+s-1}{2}}$ and $\mathsf{d}^0_{\frac{s-1}{2}}$, respectively, which are dimensions of simple modules $\mathcal{X}_j$ over $\mathcal{TL}_{\mathfrak{q},N}$

$$\begin{aligned}\dim(\mathcal{X}_j) \equiv \mathsf{d}^0_j &= \sum_{n\ge 0} \mathsf{d}_{j+np} - \sum_{n\ge t(j)+1} \mathsf{d}_{j+np-1-2(j \bmod p)} \\ &= \sum_{\substack{j'\ge j,\,(j'-j) \bmod p=0,\\ -2(j \bmod p)-1}} (-1)^{(j'-j) \bmod p}\, \mathsf{d}_{j'}, \qquad j \bmod p \ne \frac{kp-1}{2}, \quad k=0,1,\end{aligned} \tag{96}$$

where we introduce numbers $\mathsf{d}_j = \binom{N}{N/2+j} - \binom{N}{N/2+j+1}$ and a step function $t(j) \equiv t$ as

$$t = \begin{cases} 1, & \text{for} \quad j \bmod p > \dfrac{p-1}{2}, \\ 0, & \text{for} \quad j \bmod p < \dfrac{p-1}{2}. \end{cases} \tag{97}$$

We assume that $\mathsf{d}^0_j = 0$ whenever $j > L$, as usually.

We next turn to an introduction of a lattice W-algebra which uses the decompositions over $U_\mathfrak{q}s\ell(2)$ but before we give explicit examples.

4.5.1. *Examples.* We now give several examples of decompositions of the XXZ spin-chains over $U_\mathfrak{q}s\ell(2)$ for $p = 2, 3, 4$.

- For $p = 2$, we have

$$\mathcal{H}_2 = \mathsf{P}_{1,1}, \qquad \mathcal{H}_3 = 2\mathsf{X}_{2,1} \oplus \mathsf{X}_{2,2}, \qquad \mathcal{H}_4 = 2\mathsf{P}_{1,1} \oplus \mathsf{P}_{1,2}, \qquad \dots$$

- For $p = 3$, we have

$$\begin{gathered}\mathcal{H}_2 = \mathsf{X}_{1,1} \oplus \mathsf{X}_{3,1}, \qquad \mathcal{H}_3 = \mathsf{X}_{2,1} \oplus \mathsf{P}_{2,1}, \qquad \mathcal{H}_4 = \mathsf{X}_{1,1} \oplus 3\mathsf{X}_{3,1} \oplus \mathsf{P}_{1,1}, \\ \mathcal{H}_5 = \mathsf{X}_{2,1} \oplus 4\mathsf{P}_{2,1} \oplus \mathsf{X}_{3,2}, \qquad \mathcal{H}_6 = \mathsf{X}_{1,1} \oplus 9\mathsf{X}_{3,1} \oplus 4\mathsf{P}_{1,1} \oplus \mathsf{P}_{2,2}, \\ \mathcal{H}_7 = \mathsf{X}_{2,1} \oplus 13\mathsf{P}_{2,1} \oplus 6\mathsf{X}_{3,2} \oplus \mathsf{P}_{1,2}, \quad \mathcal{H}_8 = \mathsf{X}_{1,1} \oplus 28\mathsf{X}_{3,1} \oplus 13\mathsf{P}_{1,1} \oplus 7\mathsf{P}_{2,2} \oplus \mathsf{X}_{3,3}, \qquad \dots\end{gathered}$$

- For $p = 4$, we have

$$\begin{gathered}\mathcal{H}_2 = \mathsf{X}_{1,1} \oplus \mathsf{X}_{3,1}, \qquad \mathcal{H}_3 = 2\mathsf{X}_{2,1} \oplus \mathsf{X}_{4,1}, \qquad \mathcal{H}_4 = 2\mathsf{X}_{1,1} \oplus 2\mathsf{X}_{3,1} \oplus \mathsf{P}_{3,1}, \\ \mathcal{H}_5 = 4\mathsf{X}_{2,1} \oplus 4\mathsf{X}_{4,1} \oplus \mathsf{P}_{2,1}, \qquad \mathcal{H}_6 = 4\mathsf{X}_{1,1} \oplus 4\mathsf{X}_{3,1} \oplus 5\mathsf{P}_{3,1} \oplus \mathsf{P}_{1,1}, \\ \mathcal{H}_7 = 8\mathsf{X}_{2,1} \oplus 14\mathsf{X}_{4,1} \oplus 6\mathsf{P}_{2,1} \oplus \mathsf{X}_{4,2}, \qquad \mathcal{H}_8 = 8\mathsf{X}_{1,1} \oplus 8\mathsf{X}_{3,1} \oplus 20\mathsf{P}_{3,1} \oplus 6\mathsf{P}_{1,1}, \oplus \mathsf{P}_{3,2} \qquad \dots\end{gathered}$$

where the multiplicities in the front of each direct summand are dimensions of irreducible modules over $\mathcal{TL}_{\mathfrak{q},N}$ – the centralizer of $U_\mathfrak{q} s\ell(2)$.

We see from the decompositions in **4.5.1** a general pattern: the simple projective module $\mathsf{X}_{p,2}$ appears first time at $N = 2p-1$ sites. This corresponds to the first time when the algebra of endomorphisms of the module $\mathcal{H}_N$ over the small quantum group $\overline{U}_\mathfrak{q} s\ell(2)$ is larger than the algebra of endomorphisms respecting the full quantum group $U_\mathfrak{q} s\ell(2)$ (we discuss how $U_\mathfrak{q} s\ell(2)$-modules are restricted to $\overline{U}_\mathfrak{q} s\ell(2)$ after (93) and (94).) Therefore, we have isomorphisms

$$\mathrm{End}_{U_\mathfrak{q} s\ell(2)}(\mathcal{H}_N) \cong \mathrm{End}_{\overline{U}_\mathfrak{q} s\ell(2)}(\mathcal{H}_N), \qquad \text{for } 1 \le N \le 2p-2, \tag{98}$$

while we have an inclusion

$$\mathrm{End}_{U_\mathfrak{q} s\ell(2)}(\mathcal{H}_N) \subset \mathrm{End}_{\overline{U}_\mathfrak{q} s\ell(2)}(\mathcal{H}_N), \qquad \text{for } N \ge 2p-1. \tag{99}$$

We recall that the Temperley–Lieb algebra $\mathcal{TL}_{\mathfrak{q},N}$ is alternatively defined as $\mathrm{End}_{U_\mathfrak{q} s\ell(2)}(\mathcal{H}_N)$.

4.6. **Definition.** *The lattice W-algebra* $\mathcal{W}_{\mathfrak{q},N}$, for $\mathfrak{q} = e^{i\pi/p}$, is the algebra of endomorphisms respecting the action of the restricted quantum group $\overline{U}_\mathfrak{q} s\ell(2)$ on the tensor-product representation $\mathcal{H}_N = \otimes_{j=1}^N \mathsf{X}_{2,1}$.

We note that the algebra $\mathcal{W}_{\mathfrak{q},2p-1}$ is a non-trivial extension of $\mathcal{TL}_{\mathfrak{q},2p-1}$. It has dimension $\dim \mathcal{TL}_{\mathfrak{q},2p-1} + 3$. Note that the modules $\mathsf{X}_{p,1}$ and $\mathsf{X}_{p,2}$ restricted to $\overline{U}_\mathfrak{q} s\ell(2)$ are X_p^+ and $2\mathsf{X}_p^-$, respectively. This means that $\mathrm{Hom}_{\overline{U}_\mathfrak{q} s\ell(2)}(\mathsf{X}_{p,2}, \mathsf{X}_{p,1}) = 0$ and therefore three additional elements/generators are endomorphisms on $\mathsf{X}_{p,2}$ and are given by the two maps e and f mixing the two copies in the $s\ell(2)$-doublet and by h restricted to this summand. The corresponding generators in $\mathcal{W}_{\mathfrak{q},2p-1}$ can be explicitly constructed using the primitive idempotent $\boldsymbol{e}_0$ given in App. **A.2** – the normalized projector ($\boldsymbol{e}_0^2 = \boldsymbol{e}_0$) on the modules $\mathsf{X}_{p,r}$ for any *even* r.

We now turn to a definition of the lattice W-algebra $\mathcal{W}_{\mathfrak{q},N}$ using generators.

4.7. **Generators via the primitive idempotent and** $s\ell(2)$**.** We consider an extension $\mathcal{W}'_{\mathfrak{q},N}$ of the Temperley–Lieb algebra $\mathcal{TL}_{\mathfrak{q},N}$ by the generators

$$\mathcal{WB}_j^+ = \mathsf{e}_{j,j+2p-2}\, \rho_{\mathfrak{q},N}(\boldsymbol{e}_0)_{j,j+2p-2}, \tag{100}$$

$$\mathcal{WB}_j^0 = \mathsf{h}_{j,j+2p-2}\, \rho_{\mathfrak{q},N}(\boldsymbol{e}_0)_{j,j+2p-2}, \tag{101}$$

$$\mathcal{WB}_j^- = \mathsf{f}_{j,j+2p-2}\, \rho_{\mathfrak{q},N}(\boldsymbol{e}_0)_{j,j+2p-2}, \tag{102}$$

where $j = 1, 2, \dots, N-2p+2$ and we introduce the notation $\mathsf{x}_{j,j+n}$ for the n-folded comultiplication for the renormalized powers e, f, and h applied between the jth and $(j+n)$th sites and acted by identity otherwise. The algebra $\mathcal{W}'_{\mathfrak{q},N}$ commutes with $\overline{U}_\mathfrak{q} s\ell(2)$

$$\big[\mathcal{WB}_j^\alpha, \rho_{\mathfrak{q},N}(\overline{U}_\mathfrak{q} s\ell(2))\big] = 0, \qquad 1 \le j \le N-2p+2, \quad \alpha \in \{0, \pm\}, \tag{103}$$

by construction: $\mathcal{WB}_j^\alpha$ acts non-trivially on $2p-1$ tensorands between jth and $(j+2p-2)$th sites and by the identity on the other tensorands, the $2p-1$ tensorands consist the module

isomorphic to the module $\mathcal{H}_{2p-1}$ and the WB_j^α are intertwiners respecting the $\overline{U}_\mathfrak{q} s\ell(2)$-action on this $\mathcal{H}_{2p-1}$ and they do not commute with the $Us\ell(2)$ action.

We could write an explicit XXZ expression for WB_j^α using the known expression for the central idempotent $\boldsymbol{e}_0$ as a polynomial in the Casimir operator, see **A.2**:

$$\Delta^n(\boldsymbol{e}_0) = \frac{1}{4\prod_{j=1}^{p-1}(2-\mathfrak{q}^j-\mathfrak{q}^{-j})^2}\big(\Delta^n(\widetilde{\boldsymbol{C}})+2\big)\prod_{j=1}^{p-1}\big(\Delta^n(\widetilde{\boldsymbol{C}})-\mathfrak{q}^j-\mathfrak{q}^{-j}\big)^2, \tag{104}$$

where

$$\Delta^n(\widetilde{\boldsymbol{C}}) = (\mathfrak{q}-\mathfrak{q}^{-1})^2\Delta^n(\mathsf{F})\Delta^n(\mathsf{E}) + \mathfrak{q}\Delta^n(\mathsf{K}) + \mathfrak{q}^{-1}\Delta^n(\mathsf{K}^{-1}). \tag{105}$$

Then, the action of $\rho_{\mathfrak{q},N}(\boldsymbol{e}_0)_{j,j+2p-2}$ is obtained using the action (83), where we appropriately shift the indices in the σ matrices and replace N by $2p-1$. We also need formulas for $(2p-2)$-folded comultiplication of the divided powers e and f applied on the spin-$\frac{1}{2}$ chain. They are similarly obtained from (84) and (85). Using all these formulas one can write the generators WB_j^α in terms of Pauli matrices, to obtain expressions similar to TL generators e_j, but they are actually very bulky and we do not give them here.

We see from the definition which involves the $s\ell(2)$ generators that for all j the generators WB_j^α satisfy the $s\ell(2)$-relations.

4.7.1. **Proposition.** *For $1 \le j \le N-2p+2$, the generators WB_j^+, WB_j^0 and WB_j^- satisfy the $s\ell(2)$ relations*

$$[WB_j^+, WB_j^-] = 2WB_j^0, \qquad [WB_j^0, WB_j^\pm] = \pm WB_j^\pm. \tag{106}$$

and they span a basis in the adjoint representation of $s\ell(2)$:

$$[\mathsf{e}, WB_j^+] = 0, \qquad [\mathsf{e}, WB_j^0] = -WB_j^+, \qquad [\mathsf{e}, WB_j^-] = 2WB_j^0, \tag{107}$$

$$[\mathsf{f}, WB_j^+] = -2WB_j^0, \qquad [\mathsf{f}, WB_j^0] = WB_j^-, \qquad [\mathsf{f}, WB_j^-] = 0. \tag{108}$$

We therefore have that there exists an action of $s\ell(2)$ on $\mathcal{W}'_{\mathfrak{q},N}$ by derivatives of the associative multiplication.

To get more relations in $\mathcal{W}'_{\mathfrak{q},N}$, we need to express $\rho_{\mathfrak{q},N}(\boldsymbol{e}_0)_{j,j+2p-2}$ in TL generators e_k with $j \le k \le j+2p-3$. We now turn to this problem.

4.8. **Generators via the $\mathfrak{q}$-symmetrizer.** We now give an alternative description of the generators using the $\mathfrak{q}$-symmetrizers S_r projecting on the representation of the maximum spin in the tensor product of spin-$\frac{1}{2}$ representations; we assume now that $\mathfrak{q}$ is generic. The intertwiners S_r are elements from $\mathcal{TL}_{\mathfrak{q},N}$ and defined by the recursion relation

$$S_{r+1} = S_r - \frac{[r]}{[r+1]} S_r e_r S_r, \tag{109}$$

where $S_1 = 1$ and e_r is the XXZ representation of the TL-generator e_r. Using (109), we easily get first ones

$$S_2 = 1 - \frac{1}{[2]} e_1,$$
$$S_3 = 1 - \frac{[2]}{[3]}(e_1 + e_2) + \frac{1}{[3]}\{e_1, e_2\}.$$

We see that at the case of a root of unity the S_r are degenerate for $r \geq p$ (because $[p] = 0$) but not all of them. Indeed, S_2 is degenerate at $\mathfrak{q} = i$ while S_3 is well-defined and $S_3 \to 1 - e_1 e_2 - e_2 e_1$. In general, the nondegenerate ones are S_{np-1}, for any $n \geq 1$. This follows from the fact that the (projective and irreducible) module $\mathsf{X}_{p,n}$ is the limit $\mathfrak{q} \to e^{i\pi/p}$ of the np-dimensional Weyl module (generically irreducible module of the spin $\frac{np-1}{2}$) and the $\mathsf{X}_{p,n}$ is the direct summand in the limit. Therefore, there exists a projector onto this direct summand and this projector for $N = 2p - 1$ is simply given by the primitive central idempotent $\boldsymbol{e}_0$ of the quantum group. We thus observe that the primitive idempotent $\boldsymbol{e}_0$ acted on $\mathcal{H}_{2p-1}$ (and used above in the definition of the generators WB_j^α) coincides with S_{2p-1}. For example, at $\mathfrak{q} = i$ the operator $S_3 = 1 - e_1 e_2 - e_2 e_1$ equals $\rho_{\mathfrak{q},N}(\boldsymbol{e}_0)$ acted on 3 sites. This observation allows us to give a more useful definition of the lattice W-algebra generators

$$WB_j^+ = \mathsf{e}_{j,j+2p-2}\, S_{2p-1}(j), \tag{110}$$
$$WB_j^0 = \mathsf{h}_{j,j+2p-2}\, S_{2p-1}(j), \tag{111}$$
$$WB_j^- = \mathsf{f}_{j,j+2p-2}\, S_{2p-1}(j), \tag{112}$$

where $S_{2p-1}(j)$ is the Jones–Wenzl projector of the TL subalgebra generated by e_k, with $j \leq k \leq j + 2p - 2$, *i.e.*, it is defined by the recursion relation

$$S_{r+1}(j) = S_r(j) - \frac{[r]}{[r+1]} S_r(j) e_{r+j-1} S_r(j). \tag{113}$$

So, on the full spin-chain, this operator is applied between the jth and $(j + 2p - 2)$th sites and acts by identity on the other tensorands.

We first note that the generators (110)-(112) of the algebra $\mathcal{W}'_{i,N}$ (at $\mathfrak{q} = i$) coincide with the ones introduced in **2.4**, where we used a representation of the wall Brauer algebra, up to a sign factor $(-1)^j$. As in the case of free fermions, the algebra $\mathcal{W}'_{\mathfrak{q},N}$ has many defining relations in addition to the $s\ell(2)$ relations. We do not see any reasons to write all of them down because they do not actually help in studying representation theory. Moreover, we believe that most of them are relations coming from a particular representation of a bigger algebra which still needs to be discovered. Among the defining relations we can easily get few simple ones. Recall that we have relations in the TL algebra $S_r e_j = e_j S_r = 0$, for any $1 \leq j \leq r - 1$. We therefore have

$$WB_k^\alpha e_j = e_j WB_k^\alpha = 0, \qquad k \leq j \leq k + 2p - 3, \quad \alpha = \pm, 0. \tag{114}$$

Finally, we believe that $\mathcal{W}'_{\mathfrak{q},N}$ generates the full centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$. This is indeed true for $p=2$ case where the definition (13)-(15) of the lattice W-algebra coincides with $\mathcal{W}'_{\mathfrak{q}=i,N}$ defined here and we have proven Thm. **B.1** about the centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$ for this $\mathfrak{q}=i$ case. We thus finish this subsection with the following very reasonable conjecture which will be proven elsewhere.

4.9. **Conjecture.** *The associative algebra $\mathcal{W}'_{\mathfrak{q},N}$ defined by the generators WB_j^α, with $1\le j\le N-2p+2$ and $\alpha\in\{0,\pm\}$, and e_k, with $1\le k\le N-1$, in the XXZ representation is isomorphic to the centralizer $\mathcal{W}_{\mathfrak{q},N}$ of $\overline{U}_\mathfrak{q} s\ell(2)$.*

We next give a short discussion of our results on representation theory of the lattice W-algebra $\mathcal{W}_{\mathfrak{q},N}$ and relations with results on the representation theory of the chiral triplet W-algebras in the limit, for $c=-2$ and $c=0$ cases.

4.10. **Representation theory of $\mathcal{W}_{\mathfrak{q},N}$.** By the definition **4.6**, the algebra $\mathcal{W}_{\mathfrak{q},N}$ is the full centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$, *i.e.*, it is defined as $\mathcal{W}_{\mathfrak{q},N}=\mathrm{End}_{\overline{U}_\mathfrak{q} s\ell(2)}(\mathcal{H}_N)$ the algebra of all operators commuting with the small quantum group $\overline{U}_\mathfrak{q} s\ell(2)$. Its representation theory can be thus studied using the decomposition of the XXZ spin-chain over $\overline{U}_\mathfrak{q} s\ell(2)$. Such a decomposition can be easily obtained by restriction of the $U_\mathfrak{q} s\ell(2)$-module $\mathcal{H}_N$ in (95) on the subalgebra $\overline{U}_\mathfrak{q} s\ell(2)$. The idea is then to describe the multiplicity spaces in front of $\overline{U}_\mathfrak{q} s\ell(2)$ direct summands. These multiplicity spaces are then irreducible modules over $\mathcal{W}_{\mathfrak{q},N}$. The next step is to describe all possible homomorphisms between the $\overline{U}_\mathfrak{q} s\ell(2)$ direct summands. These homomorphisms represent the action of $\mathcal{W}_{\mathfrak{q},N}$ in its reducible and indecomposable modules. We only announce our results about $\mathcal{W}_{\mathfrak{q},N}$-modules in order to make the paper not too long. All the necessary details will be given in a forthcoming publication [37].

We begin with description of projective covers for simple $\mathcal{W}_{\mathfrak{q},N}$-modules and describe then their content with respect to the TL subalgebra. Note that $\mathcal{W}_{\mathfrak{q},N}$ is represented faithfully in the spin chain, by its definition as the subalgebra in the algebra of all operators acting on $\mathcal{H}_N$. Therefore, projective covers of irreducible modules can be identified with indecomposable submodules in $\mathcal{H}_N$ of maximum dimension.

The algebra $\mathcal{W}_{\mathfrak{q},N}$ has the following projective covers. The modules $\mathcal{X}_p^\pm$ are irreducible and projective simultaneously. There are $p-1$ cells (taking into account odd and even values of N) where reducible but indecomposable modules exist. Each cell contains projective modules $\mathcal{P}_s^+$, $\mathcal{P}_s^-$ and $\mathcal{P}_s^0$ with the subquotient structures given in Fig. 1 (from the left to the right). These modules are projective covers of irreducible modules denoted by $\mathcal{X}_s^+$, $\mathcal{X}_{p-s}^-$, and $\mathcal{Y}_s$, respectively. The decomposition of the irreducible $\mathcal{W}_{\mathfrak{q},N}$-modules over the two commuting algebras, $\mathcal{TL}_{\mathfrak{q},N}$ and $Us\ell(2)$, can be written as

$$\mathcal{Y}_s=\mathcal{X}_{\frac{s-1}{2}}\otimes\mathbb{C}^1,\qquad 1\le s\le p-1, \tag{115}$$

$$\mathcal{X}_s^+=\bigoplus_{r\ge1}\mathcal{X}_{\frac{2rp-s-1}{2}}\otimes\mathbb{C}^{2r-1},\qquad 1\le s\le p, \tag{116}$$




FIGURE 1. Projective covers of irreducible $\mathcal{W}_{\mathfrak{q},N}$-modules denoted by $\mathcal{X}_s^+$, $\mathcal{X}_{p-s}^-$, and $\mathcal{Y}_s$. Same diagrams also describe subquotient structure of indecomposable modules over the chiral triplet W-algebra from [4].

$$\mathcal{X}_{p-s}^- = \bigoplus_{r\geq 1} \mathcal{X}_{\frac{2rp+s-1}{2}} \otimes \mathbb{C}^{2r}, \qquad 0 \leq s \leq p-1, \tag{117}$$

where the direct sums are finite and as usually we imply that $\mathcal{X}_{j>N} \equiv 0$ and $s+N = 1 \bmod 2$; the tensorands $\mathbb{C}^r$ denote the r-dimensional modules over $s\ell(2)$. We note that the modules $\mathcal{X}_{p-s}^-$ are non-zero only for $N \geq 2p-1$. For $N < 2p-1$, we have modules $\mathcal{Y}_s$ and $\mathcal{X}_s^+$ with trivial $Us\ell(2)$ content. They are essentially the irreducible modules over $\mathcal{TL}_{\mathfrak{q},N}$, in accordance with (98).

The dimensions of irreducible $\mathcal{W}_{\mathfrak{q},N}$-modules can be expressed in terms of the multiplicities d_j^0 introduced above in (96). The dimension of $\mathcal{Y}_s$ is equal to $\mathsf{d}^0_{\frac{s-1}{2}}$ and the module is the singlet with respect to the $Us\ell(2)$ action. The dimensions of the irreducible $\mathcal{W}_{\mathfrak{q},N}$-modules $\mathcal{X}_s^\pm$ are

$$\dim(\mathcal{X}_s^+) = \sum_{r\geq 1}(2r-1)\mathsf{d}^0_{\frac{2rp-s-1}{2}}, \qquad \dim(\mathcal{X}_s^-) = \sum_{r\geq 1} 2r\, \mathsf{d}^0_{\frac{(2r+1)p-s-1}{2}}.$$

Finally, we give the $\mathcal{TL}_{\mathfrak{q},N} \otimes Us\ell(2)$ content of the projective $\mathcal{W}_{\mathfrak{q},N}$-modules

$$\begin{aligned} \mathcal{P}_s^+|_{\mathcal{TL}_{\mathfrak{q},N}\otimes Us\ell(2)} &= \bigoplus_{n\geq 1} \mathcal{P}_{\frac{2np-s-1}{2}} \otimes \mathbb{C}^{2n-1}, \qquad 1 \leq s \leq p, \\ \mathcal{P}_{p-s}^-|_{\mathcal{TL}_{\mathfrak{q},N}\otimes Us\ell(2)} &= \bigoplus_{n\geq 1} \mathcal{P}_{\frac{2np+s-1}{2}} \otimes \mathbb{C}^{2n} \qquad 0 \leq s \leq p-1, \end{aligned} \qquad s+N = 1 \bmod 2, \tag{118}$$

where the direct sums are finite of course, similarly to those for the simple modules above, and the subquotient structure of the projective $\mathcal{TL}_{\mathfrak{q},N}$-modules $\mathcal{P}_j$ can be found, *e.g.*, in [32] and we use the identity $\mathcal{P}_{\frac{rp-1}{2}} = \mathcal{X}_{\frac{rp-1}{2}}$ for modules over $\mathcal{TL}_{\mathfrak{q},N}$.

We note also that the modules $\mathcal{P}_{p-s}^-$ do not exist unless $N \geq 2p-1$. For $N < 2p-1$, we have projective modules $\mathcal{P}_s^0$ and $\mathcal{P}_s^+$ with trivial $Us\ell(2)$ content – the two subquotients in the first diagram on the left in Fig. 1 are actually absent (because of our notation $\mathcal{X}_{j>N} = 0$ for $\mathcal{TL}_{\mathfrak{q},N}$-modules), and a module $\mathcal{P}_s^+$ has only three irreducible subquotients. These projectives

are essentially the projective modules over $\mathcal{TL}_{\mathfrak{q},N}$ and the algebra $\mathcal{W}_{\mathfrak{q},N}$ has thus the same representation theory whenever $N < 2p-1$, in accordance with (98).

Finally, the decomposition of the full spin-chain over $\mathcal{W}_{\mathfrak{q},N}$ can be written as

$$\mathcal{H}_N|_{\mathcal{W}_{\mathfrak{q},N}} \cong \bigoplus_{\substack{s=N \bmod 2+1,\\ s+N=1 \bmod 2}}^{p-1} \left(s\mathcal{P}_s^+ \oplus (p-s)\mathcal{P}_{p-s}^-\right) \oplus \delta_{(p+N) \bmod 2,1}\, p\,\mathcal{P}_p^+ \oplus \delta_{N \bmod 2,1}\, p\,\mathcal{P}_p^-, \tag{119}$$

where $\delta_{a,b}$ is the usual Kronecker symbol.

4.11. **Two scaling limits: $(p-1,p)$ and $(1,p)$ theories.** Recall that in the free fermions case we already took the scaling limit of our spin-chains, and thus of the lattice W-algebra projective modules. It is easy to see that in this case ($p=2$) the modules $\mathcal{Y}_s$ are absent and we have only two reducible and indecomposable projectives $\mathcal{P}_1^\pm$ of the "diamond" structure (four and not five subquotients) on even number of sites. For odd number N, the spin-chain is semisimple and both the projectives $\mathcal{P}_2^\pm$ are irreducible. This subquotient structure persists in the limit and give exactly the modules over the triplet W-algebra [11], with the same character of course. This result is not very surprising as far as we consider free theories.

It is more interesting to consider the case $p=3$ or $\mathfrak{q}=e^{\frac{i\pi}{3}}$ which corresponds in the scaling limit to a $c=0$ logarithmic CFT [3]. Indeed, the generating function of energy levels of the Hamiltonian $H(\mathfrak{q})$ from (78) is well known [23] thanks to the Bethe ansatz, and the partition function for scaling states in the XXZ spin-chain can be written as

$$\lim_{N\to\infty} \sum_{\text{states } i} q^{\frac{N}{2\pi v_F}(E_i(N)-Ne_\infty)} = q^{-c/24} \sum_{j\geq 0} (2j+1) \frac{q^{h_{1,1+2j}} - q^{h_{1,-1-2j}}}{\prod_{n=1}^\infty (1-q^n)}, \tag{120}$$

where $v_F = \frac{\pi \sin\gamma}{\gamma}$ (with $2\cos\gamma = \mathfrak{q}+\mathfrak{q}^{-1}$) is the Fermi velocity, with the central charge

$$c = c_{p-1,p} = 1 - \frac{6}{p(p-1)}, \tag{121}$$

and $E_i(N)$ is the eigenvalue of the i^{th} (counted from the vacuum) eigenstate of $H = -\sum_i e_i$. Here, we have also subtracted the density $e_\infty = \lim_{N\to\infty} E_0(N)/N$, with $E_0(N)$ the ground-state energy. The expression at the multiplicity $(2j+1)$ on the right-hand side of (120) coincides with the Virasoro character $\mathrm{Tr}\, q^{L_0-c/24}$ of the so-called *Kac module* with conformal weight $h_{1,1+2j}$ defined as a quotient of the Verma module as $\mathcal{V}_{h_{1,1+2j}}/\mathcal{V}_{h_{1,-1-2j}}$ by the submodule corresponding to the singular vector at level $2j+1$. We use here the standard notation for the conformal weight

$$h_{r,s} = \frac{(pr-(p-1)s)^2 - 1}{4p(p-1)} \tag{122}$$

and denote by $\mathcal{V}_h$ the Virasoro Verma module generated from the highest-weight state of weight h.

It is also well-established [3] (see also [32]) that simple TL modules $\mathcal{X}_j$ go over in the scaling limit to simple Virasoro modules with the highest weight $h_{1,2j+1}$, which we will simply denote

by $\mathcal{X}_{1,2j+1}$. Then, using the decomposition (115) of the simple $\mathcal{Y}_s$ modules, we see that they correspond to the Virasoro irreducible representations from the $(p-1,p)$ minimal model Kac table. In the case $p=3$ or $c=0$, we have only one module $\mathcal{Y}_1$ which has dimension 1 for any N and this module correspond to the unique operator (identity) in the $(2,3)$ Kac table. For any $p\geq 2$, these are also simple modules over the triplet chiral W-algebra first observed in [4].

Using now the decomposition (116) and (117) of the simple $\mathcal{X}_s^{\pm}$ modules and the identification of the simple TL modules with simple Virasoro modules in the scaling limit, we can compute the characters of their limits using the decomposition over the Virasoro algebra and $Us\ell(2)$ (we use the same notation for the limits of $\mathcal{W}_{\mathfrak{q},N}$-modules):

$$\mathcal{X}_s^{+}=\bigoplus_{r\geq 1}\mathcal{X}_{1,2rp-s}\otimes\mathbb{C}^{2r-1},\qquad 1\leq s\leq p, \tag{123}$$

$$\mathcal{X}_{p-s}^{-}=\bigoplus_{r\geq 1}\mathcal{X}_{1,2rp+s}\otimes\mathbb{C}^{2r},\qquad 0\leq s\leq p-1, \tag{124}$$

where now the direct sums are infinite and are taken over all $s\ell(2)$-spins of a fixed parity; we also recall that $\mathcal{X}_{1,j}$ is the simple Virasoro module of the weight $h_{1,j}$. The characters of $\mathcal{X}_s^{\pm}$ are then expressed exactly as in [4].

The structure of projective modules $\mathcal{P}_s^{\pm}$ does not depend on the number of sites and should thus persists in the scaling limit. We thus get the same diagrams in the scaling limit as those in Fig. 1. It is interesting to note that indecomposable modules over the chiral triplet W-algebra for $c=0$ ($p=3$) with the same subquotient structure were proposed in [36] (note also that for any $(p-1,p)$ theory similar modules with 5, 4, and 2 subquotients involving the minimal model content were constructed in [4]).

It is interesting that we can take another scaling limit using the same algebras. Recall that the Hamiltonian H spectrum has two points of condensation of energy levels [33] – the vacuum of $H(\mathfrak{q})$ and the vacuum of $-H(\mathfrak{q})$. The low-lying spectrum of $-H(\mathfrak{q})$ is the same as for the Hamiltonian $H(-\mathfrak{q}^{-1})$ which has the Heisenberg coupling with the opposite sign. Indeed their spectra are related by the similarity transformation

$$-H(\mathfrak{q})=e^{-\mathrm{i}\pi\sum_{j=1}^{N}j\sigma_j^z}H(-\mathfrak{q}^{-1})e^{\mathrm{i}\pi\sum_{j=1}^{N}j\sigma_j^z}. \tag{125}$$

This observation tells us that the generating function of low-lying states for $-H(\mathfrak{q})$ converges to characters from the $(1,p)$ theories. Indeed note that $-\mathfrak{q}^{-1}=\mathfrak{q}^{p-1}$ and recall that the generating function of low-lying states for $H\big(e^{\frac{i\pi(p-p')}{p}}\big)$ converges to characters from the (p',p) theory [23]. Then, at this limit the structure of the projective covers will be slightly different – the subquotients $\mathcal{Y}_s$ corresponding to the minimal model content disappear in the $(1,p)$ scaling limit. It is indeed easy to see at the case $p=3$ when we have just one $\mathcal{Y}_{s=1}$, and it is spanned by one state which is now highest-energy state for the Hamiltonian $-H(\mathfrak{q})$ (or $H(-\mathfrak{q}^{-1})$). This state therefore will go to "infinity" in the limit. We expect that similarly all

subquotients $\mathcal{Y}_s$ have just one point of condensation of energy levels and they thus disappear in the second, $(1,p)$, scaling limit. So, all the diagrams of the reducible but indecomposable triplet W-algebra projectives will have only 4 subquotients in the limit corresponding to the $(1,p)$ theories, as it is expected.

5. Conclusions

We believe we have reached our main goal of understanding the lattice origin and analog of W-algebra symmetry in chiral (boundary) LCFTs. The next step should be the analysis of what happens in the non chiral case, that is for bulk LCFTs. There is now a lot of evidence that the bulk boundary relationship is extremely complicated in the logarithmic case, and that naive "periodicized" versions of the XXZ spin chain or the underlying loop models will not have W-symmetry. What has to be done to restore this symmetry then remains an open question of crucial importance, and we think the study of lattice W-algebras in this context will be the source of further progress.

The case $c=-2$ has also shed light on the mathematical nature of the relationship between the Temperley-Lieb and the Virasoro algebras, or the lattice and continuum W-algebras. For a finite chain, the Fourier modes H_n^r and $W_n^{\alpha,r}$ define a finite dimensional algebra whose structure constants are function of the size N. The scaling limit involves not only taking $N\to\infty$ but also normal ordering with respect to the ground state. In this scaling limit, the generators H_n^r and $W_n^{\alpha,r}$ expand on elements from the enveloping algebra of the Virasoro and the triplet W-algebra. Infinitely many lattice "approximations" converge to the same limit in leading order – *e.g.*, all the H_n^r go to $(-1)^r L_n + L_{-n}$ – but they differ at next to leading order. While our calculations were considerably simplified by the underlying presence of free fermions, we believe it should be possible to extend the analysis to $\mathfrak{q}$ another root of unity. It is important to note that different Hamiltonians will lead, for the same lattice algebra, to different continuum limits, as exemplified above in the $(1,p)$ and $(p-1,p)$ cases.

To finish, we now would like to get back to the algebra of zero modes in the $\mathfrak{q}=i$ case. The operators H_0^r and $W_0^{\alpha,r}$ have the following properties

$$[H_0^r, H_0^s]=0, \qquad [H_0^r, W_0^{\alpha,s}]=0 \tag{126}$$

and

$$[W_0^{0,r}, W_0^{+,s}] = -8W_0^{+,r+s+2} + 8W_0^{+,r+s}, \tag{127}$$

$$[W_0^{0,r}, W_0^{-,s}] = 8W_0^{-,r+s+2} - 8W_0^{-,r+s}, \tag{128}$$

$$[W_0^{+,r}, W_0^{-,s}] = 4W_0^{0,r+s+2} - 4W_0^{0,r+s}, \tag{129}$$

for $r,s\in 2\mathbb{N}_0$. For finite N operators H_0^r and $W_0^{\alpha,r}$ are linearly dependent

$$H_0^N = \sum_{r=0}^{\frac{N}{2}-1} (-1)^{\frac{N}{2}-r-1} S_{\frac{N}{2}-r}\big(\cos^2\pi\frac{0}{N}, \cos^2\pi\frac{1}{N}, \cos^2\pi\frac{2}{N}, \dots, \cos^2\pi\frac{N/2-1}{N}\big) H_0^{2r}, \tag{130}$$

$$W_0^{\alpha,N-2} = \sum_{r=0}^{\frac{N}{2}-2} (-1)^{\frac{N}{2}-r} S_{\frac{N}{2}-r-1}\big(\cos^2\pi\frac{1}{N}, \cos^2\pi\frac{2}{N}, \dots, \cos^2\pi\frac{N/2-1}{N}\big) W_0^{\alpha,2r} \tag{131}$$

where $S_r(x_1, x_2, \dots x_k)$ is the rth elementary symmetric polynomial in the variables $x_1, x_2, \dots x_k$. We also recall that H_0^0 coincides with the Hamiltonian H, thus we have $4N-10$ currents that commute with the Hamiltonian.

Interestingly, the algebra of $W_0^{\alpha,r}$ can be identified through a quotient algebra of the $s\ell(2)$ loop algebra in the following way

$$W_0^{+,r} = 4(t^2-1)t^r e, \qquad W_0^{-,r} = -4(t^2-1)t^r f, \qquad W_0^{0,r} = -8(t^2-1)t^r h, \tag{132}$$

where the parameter t satisfies the equation

$$P(t) \equiv (t^2-1)\sum_{r=0}^{\frac{N}{2}-1} (-1)^r S_r\big(\cos^2\pi\frac{1}{N}, \cos^2\pi\frac{2}{N}, \dots, \cos^2\pi\frac{N/2-1}{N}\big) t^{N-2r-2} = 0. \tag{133}$$

Recalling the well known property of elementary symmetric polynomials

$$\prod_{j=1}^{n}(t-t_j) = \sum_{r=0}^{n}(-1)^r S_r(t_1, \dots, t_n) t^{n-r},$$

we obtain that the solutions of the equation (133) are $\{\pm\cos\frac{\pi j}{N},\ 0 \le j \le N/2-1\}$. We have thus the representation of the loop algebra $s\ell(2)\otimes\mathbb{C}[t,t^{-1}]$ realized by the quotient by the ideal generated by the polynomial $P(t)$ from (133), *i.e.*, we have relations $P(t)e = 0$, $P(t)f = 0$, *etc.* It is obvious that this representation (or the quotient) of the $s\ell(2)$ loop algebra is isomorphic to the following tensor-product representation

$$V[t_1, \dots, t_N] \equiv \bigotimes_{j=0}^{N/2-1} \big(\mathbb{C}^2\big[\cos(\pi j/N)\big] \otimes \mathbb{C}^2\big[-\cos(\pi j/N)\big]\big), \tag{134}$$

where by $\mathbb{C}[t_j]$ we denote the so-called evaluation representation of the loop algebra – any element $p(t)x$ acts by $p(t_j)x$, with $p(t)\in\mathbb{C}[t,t^{-1}]$ and $x\in s\ell(2)$. The action on the full spin chain (134) (defined on N sites) is then given by the usual repeated comultiplication of a Lie algebra: $\Delta(x) = x\otimes\mathbf{1} + \mathbf{1}\otimes x$.

Therefore, the zero modes of the $W^{\alpha,r}$ generators, with $\alpha\in\{\pm,0\}$, can be written as

$$W_0^{\alpha,r} = \sum_{j=1}^{N}(t_j^2-1)t_j^r\, x_\alpha, \qquad x_+ = 4e,\ x_0 = -8h,\ x_- = -4f, \tag{135}$$

where we also assume that all t_j's are distinct and are from the set $\{\pm\cos\frac{\pi j}{N},\ 0\le j\le N/2-1\}$. Note that this expression is given in a different basis than the usual spin-basis of the XX chains.

We note further that the representation (134) of the loop algebra is irreducible because all the evaluation parameters t_j are distinct, see *e.g.* [34]. For the subalgebra generated

by the zero modes $W_0^{\alpha,r}$, it is a reducible representation. Indeed the representation (134) is isomorphic to the initial spin-chain representation where the zero modes are represented by (52)-(54) and the former one is reducible even for the full lattice W-algebra $\mathcal{W}_{i,N}$ – due to presence of the fermionic zero modes and the decomposition onto bosonic and fermionic states. The spin-chain representation of $\mathcal{W}_{i,N}$ is a direct sum of two reducible but indecomposable representations, as discussed earlier in the more general context of XXZ spin-chains. It is easy to see that the indecomposability here is due the action of non-zero modes $W_n^{\alpha,r}$, as they contain fermionic zero modes $\eta_0^{\pm}$, while the action of $W_0^{\alpha,r}$ is semisimple. The module $V[t_1,\dots,t_N]$ over the algebra of zero modes is thus semisimple and it is decomposed as (for even N)

$$V[t_1,\dots,t_N]|_{\langle W_0^{\alpha,r}\rangle} = 4\big(\mathcal{X}^+ \oplus \mathcal{X}^-\big),$$

where $\mathcal{X}^{\pm}$ are irreducible $\mathcal{W}_{i,N}$-modules and they consist of bosonic and fermionic states, respectively, generated from the vacuum state by negative fermionic modes $\eta_{n<0}^{\pm}$. Each module $\mathcal{X}^{\pm}$ has dimension 2^{N-3}. We have a similar decomposition for odd N but with multiplicity 2 instead of 4.

The algebra of the zero modes $W_0^{\alpha,r}$ is reminiscent of the Onsager algebra [31] that appears in Onsager's original solution of the Ising model, or the chiral Potts model. Like our algebra of zero modes, the Onsager algebra is also identified as a subalgebra of the $s\ell(2)$ loop algebra [35], but a different subalgebra from the one we have found here. Now the use of such an identification in the context of the Ising or chiral Potts model is that it leads to the spectrum of the Hamiltonian, which in these cases is a particular element of the loop algebra. Things are a bit different for the XX spin chain we are considering. The boundary terms prevent the usual analysis using the Onsager algebra. Meanwhile, we have another "version" of this algebra, where the Hamiltonians H_0^r are somehow "disconnected", since they commute with all the zero modes. Of course, we can decide to compute the spectrum of the conserved quantities $W_0^{0,r}$ nonetheless. Indeed using the identification (135) we obtain

$$\mathrm{Spec}(W_0^{0,r}) = \Big\{\sum_{j=1}^{N} m_j(t_j^2-1)t_j^r;\ m_j=\pm 4,\ t_j^2=\cos\frac{\pi j}{N},\ 0\le j\le \frac{N}{2}-1\Big\}. \tag{136}$$

where N is the number of sites. Of course, this spectrum (the set of eigenvalues of $W_0^{0,r}$) coincides with the eigenvalues extracted from the fermionic expression (53) at $n=0$.

It is natural to wonder at this stage whether higher spin representations, or different choices of the parameters t_j, are related with interesting new models exhibiting some sort of W-algebra symmetry. Another question is whether the lattice W-algebras for other roots of unity have anything to do with the $s\ell(2)$ loop algebra.

A last remark of interest concerns the fact that the zero modes H_0^r and $W_0^{\alpha,s}$ commute, which implies that our XX spin chain has more conserved quantities than the H_0^r, which are the standard quantities obtainable from derivatives of the transfer matrix with respect to the spectral parameter. Of course, since we are dealing with free fermions, the existence of

other conserved quantities apart from the H_0^r is obvious. We have not been able however to extend the commutation of the zero modes to other values of $\mathfrak{q}$. It is tempting to speculate that there are other possible lattice models for which such additional conserved quantities could be built, and that they would be better candidates to reproduce the bulk W-symmetric LCFTs in the continuum limit.

Acknowledgments: We thank B.L. Feigin, D. Ridout, V. Schomerus and A.M. Semikhatov for discussions. This work was supported in part by the ANR Project 2010 Blanc SIMI 4: DIME. The work of AMG was supported in part by Marie-Curie IIF fellowship, the RFBR grant 10-01-00408 and the RFBR–CNRS grant 09-01-93105. The work of IYuT is partially supported by the grant RFBR 11-02-00685.

Appendix A. Lusztig quantum group $U_{\mathfrak{q}}s\ell(2)$

Here, we collect standard expressions for the quantum group $U_{\mathfrak{q}}s\ell(2)$ we use in the definition of the lattice version of the triplet W-algebra $\mathcal{W}_{\mathfrak{q},N}$, with $\mathfrak{q}=e^{i\pi/p}$ and $p\geq 2$. We introduce standard notation for $\mathfrak{q}$-numbers $[n]=\dfrac{\mathfrak{q}^n-\mathfrak{q}^{-n}}{\mathfrak{q}-\mathfrak{q}^{-1}}$ and set $[n]!=[1][2]\dots[n]$.

A.1. **Defining relations.** The Lusztig quantum group $U_{\mathfrak{q}}s\ell(2)$ with $\mathfrak{q}=e^{i\pi/p}$, for any integer $p\geq 2$, is generated by E, F, and K satisfying the standard relations for the quantum $s\ell(2)$,

$$\mathsf{K}\mathsf{E}\mathsf{K}^{-1}=\mathfrak{q}^2\mathsf{E},\quad \mathsf{K}\mathsf{F}\mathsf{K}^{-1}=\mathfrak{q}^{-2}\mathsf{F},\quad [\mathsf{E},\mathsf{F}]=\frac{\mathsf{K}-\mathsf{K}^{-1}}{\mathfrak{q}-\mathfrak{q}^{-1}}, \tag{137}$$

with the relations,

$$\mathsf{E}^p=\mathsf{F}^p=0,\quad \mathsf{K}^{2p}=\mathbf{1},$$

and additionally by the divided powers $\mathsf{f}\sim\mathsf{F}^p/[p]!$ and $\mathsf{e}\sim\mathsf{E}^p/[p]!$ which satisfy the usual $s\ell(2)$-relations:

$$[\mathsf{h},\mathsf{e}]=\mathsf{e},\qquad [\mathsf{h},\mathsf{f}]=-\mathsf{f},\qquad [\mathsf{e},\mathsf{f}]=2\mathsf{h}.$$

There are also 'mixed' relations

$$[\mathsf{h},\mathsf{K}]=0,\qquad [\mathsf{E},\mathsf{e}]=0,\qquad [\mathsf{K},\mathsf{e}]=0,\qquad [\mathsf{F},\mathsf{f}]=0,\qquad [\mathsf{K},\mathsf{f}]=0, \tag{138}$$

$$[\mathsf{F},\mathsf{e}]=\frac{1}{[p-1]!}\mathsf{K}^p\frac{\mathfrak{q}\mathsf{K}-\mathfrak{q}^{-1}\mathsf{K}^{-1}}{\mathfrak{q}-\mathfrak{q}^{-1}}\mathsf{E}^{p-1},\qquad [\mathsf{E},\mathsf{f}]=\frac{(-1)^{p+1}}{[p-1]!}\mathsf{F}^{p-1}\frac{\mathfrak{q}\mathsf{K}-\mathfrak{q}^{-1}\mathsf{K}^{-1}}{\mathfrak{q}-\mathfrak{q}^{-1}}, \tag{139}$$

$$[\mathsf{h},\mathsf{E}]=\frac{1}{2}\mathsf{E}A,\quad [\mathsf{h},\mathsf{F}]=-\frac{1}{2}A\mathsf{F}, \tag{140}$$

where

$$A=\sum_{s=1}^{p-1}\frac{(u_s(\mathfrak{q}^{-s-1})-u_s(\mathfrak{q}^{s-1}))\mathsf{K}+\mathfrak{q}^{s-1}u_s(\mathfrak{q}^{s-1})-\mathfrak{q}^{-s-1}u_s(\mathfrak{q}^{-s-1})}{(\mathfrak{q}^{s-1}-\mathfrak{q}^{-s-1})u_s(\mathfrak{q}^{-s-1})u_s(\mathfrak{q}^{s-1})}\,u_s(\mathsf{K})\boldsymbol{e}_s \tag{141}$$

with the polynomials $u_s(\mathsf{K}) = \prod_{n=1,\, n\neq s}^{p-1}(\mathsf{K}-\mathfrak{q}^{s-1-2n})$, and $\boldsymbol{e}_s$ are central primitive idempotents defined just below following [26]. The relations (137)-(141) are the defining relations of the quantum group $U_\mathfrak{q} s\ell(2)$.

A.2. **Central idempotents.** We recall the primitive central idempotents in $\overline{U}_\mathfrak{q} s\ell(2)$ [26]:

$$\boldsymbol{e}_s = \frac{1}{\psi_s(\beta_s)}\big(\psi_s(\widetilde{\boldsymbol{C}}) - \frac{\psi_s'(\beta_s)}{\psi_s(\beta_s)}(\widetilde{\boldsymbol{C}}-\beta_s)\psi_s(\widetilde{\boldsymbol{C}})\big), \quad 1\le s\le p-1,$$

$$\boldsymbol{e}_0 = \frac{1}{\psi_0(\beta_0)}\psi_0(\widetilde{\boldsymbol{C}}), \qquad \boldsymbol{e}_p = \frac{1}{\psi_p(\beta_p)}\psi_p(\widetilde{\boldsymbol{C}}),$$

with the polynomials

$$\psi_s(x) = (x-\beta_0)\,(x-\beta_p)\prod_{\substack{j=1\\ j\neq s}}^{p-1}(x-\beta_j)^2, \quad 1\le s\le p-1,$$

$$\psi_0(x) = (x-\beta_p)\prod_{j=1}^{p-1}(x-\beta_j)^2, \quad \psi_p(x) = (x-\beta_0)\prod_{j=1}^{p-1}(x-\beta_j)^2,$$

where $\beta_j = \mathfrak{q}^j + \mathfrak{q}^{-j}$, and the Casimir element is

$$\widetilde{\boldsymbol{C}} = (\mathfrak{q}-\mathfrak{q}^{-1})^2\mathsf{EF} + \mathfrak{q}^{-1}\mathsf{K} + \mathfrak{q}\mathsf{K}^{-1}.$$

A.3. **Comultiplication in $U_\mathfrak{q} s\ell(2)$.** The quantum group $U_\mathfrak{q} s\ell(2)$ has the Hopf-algebra structure with the comultiplication

$$\Delta(\mathsf{E}) = \mathbf{1}\otimes\mathsf{E} + \mathsf{E}\otimes\mathsf{K}, \quad \Delta(\mathsf{F}) = \mathsf{K}^{-1}\otimes\mathsf{F} + \mathsf{F}\otimes\mathbf{1}, \quad \Delta(\mathsf{K}) = \mathsf{K}\otimes\mathsf{K}, \tag{142}$$

$$\Delta(\mathsf{e}) = \mathsf{e}\otimes\mathbf{1} + \mathsf{K}^p\otimes\mathsf{e} + \frac{1}{[p-1]!}\sum_{r=1}^{p-1}\frac{\mathfrak{q}^{r(p-r)}}{[r]}\mathsf{K}^p\mathsf{E}^{p-r}\otimes\mathsf{E}^r\mathsf{K}^{-r}, \tag{143}$$

$$\Delta(\mathsf{f}) = \mathsf{f}\otimes\mathbf{1} + \mathsf{K}^p\otimes\mathsf{f} + \frac{(-1)^p}{[p-1]!}\sum_{s=1}^{p-1}\frac{\mathfrak{q}^{-s(p-s)}}{[s]}\mathsf{K}^{p+s}\mathsf{F}^s\otimes\mathsf{F}^{p-s}. \tag{144}$$

The antipode and counity are not used in the paper but a reader can find them, for exmaple, in [25].

The N-folded comultiplication for the capital generators E and F is

$$\begin{aligned}\Delta^N(\mathsf{E}) &= \sum_{j=1}^{N+1}\underbrace{\mathbf{1}\otimes\ldots\otimes\mathbf{1}}_{j-1}\otimes\mathsf{E}\otimes\mathsf{K}\otimes\ldots\otimes\mathsf{K},\\ \Delta^N(\mathsf{F}) &= \sum_{j=1}^{N+1}\underbrace{\mathsf{K}^{-1}\otimes\ldots\otimes\mathsf{K}^{-1}}_{j-1}\otimes\mathsf{F}\otimes\mathbf{1}\otimes\ldots\otimes\mathbf{1},\end{aligned} \tag{145}$$

and for the divided powers e and f is

$$\Delta^N(\mathsf{e}) = \sum_{n=1}^{N+1}\bigotimes_{j=1}^{n-1}\mathsf{K}^p\otimes\mathsf{e}\otimes\bigotimes_{j=n+1}^{N+1}\mathbf{1} \tag{146}$$

$$+\sum_{\substack{0\le j_1,j_2,\dots,j_{N+1}\le p-1\\ j_1+j_2+\dots+j_{N+1}=p}} \frac{\mathfrak{q}^{\sum_{r<s} j_r j_s}}{[j_1]![j_2]!\dots[j_{N+1}]!} \bigotimes_{m=1}^{N+1} \mathsf{E}^{j_m}\mathsf{K}^{p+\sum_{s=1}^{m-1} j_s},$$

$$\Delta^N(\mathsf{f}) = \sum_{n=1}^{N+1} \bigotimes_{j=1}^{n-1} \mathsf{K}^p \otimes \mathsf{f} \otimes \bigotimes_{j=n+1}^{N+1} \mathbf{1} \tag{147}$$

$$+(-1)^{p-1} \sum_{\substack{0\le j_1,j_2,\dots,j_{N+1}\le p-1\\ j_1+j_2+\dots+j_{N+1}=p}} \frac{\mathfrak{q}^{\sum_{r<s} j_r j_s}}{[j_1]![j_2]!\dots[j_{N+1}]!} \bigotimes_{m=1}^{N+1} \mathsf{F}^{j_m}\mathsf{K}^{p+\sum_{s=1}^{m} j_s}.$$

A.4. **Standard spin-chain notations.** We note the Hopf-algebra homomorphism

$$\mathsf{E} \mapsto S^+ k, \qquad \mathsf{F} \mapsto k^{-1} S^-, \qquad \text{with} \quad k = \sqrt{\mathsf{K}},$$

where we introduced the more usual (in the spin-chain literature [23, 30][2]) quantum group generators

$$S^\pm = \sum_{1\le j\le N} \mathfrak{q}^{-\sigma_1^z/2} \otimes \dots \mathfrak{q}^{-\sigma_{j-1}^z/2} \otimes \sigma_j^\pm \otimes \mathfrak{q}^{\sigma_{j+1}^z/2} \otimes \dots \otimes \mathfrak{q}^{\sigma_N^z/2} \tag{148}$$

together with $k = \mathfrak{q}^{S^z}$ and the relations

$$kS^\pm k^{-1} = \mathfrak{q}^{\pm 1} S^\pm, \qquad \left[S^+, S^-\right] = \frac{k^2 - k^{-2}}{\mathfrak{q} - \mathfrak{q}^{-1}},$$
$$\Delta(S^\pm) = k^{-1} \otimes S^\pm + S^\pm \otimes k.$$

A.4.1. *The case of XX spin-chains.* For $p = 2$ or "XX spin-chain" case, the $(N-1)$-folded coproduct of the renormalized powers e and f reads

$$\Delta^{N-1}\mathsf{e} = \sum_{j=1}^{N} \underbrace{\mathbf{1} \otimes \dots \otimes \mathbf{1}}_{j-1} \otimes \mathsf{e} \otimes \mathsf{K}^2 \otimes \dots \otimes \mathsf{K}^2 + \tag{149}$$
$$+ \mathfrak{q} \sum_{t=0}^{N-2} \sum_{j=1}^{N-1-t} \underbrace{\mathbf{1} \otimes \dots \otimes \mathbf{1}}_{j-1} \otimes \mathsf{E} \otimes \underbrace{\mathsf{K} \otimes \dots \otimes \mathsf{K}}_{t} \otimes \mathsf{EK} \otimes \mathsf{K}^2 \otimes \dots \otimes \mathsf{K}^2$$

and

$$\Delta^{N-1}\mathsf{f} = \sum_{j=1}^{N} \underbrace{\mathsf{K}^2 \otimes \dots \otimes \mathsf{K}^2}_{j-1} \otimes \mathsf{f} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1} + \tag{150}$$
$$+ \mathfrak{q}^{-1} \sum_{t=0}^{N-2} \sum_{j=1}^{N-1-t} \underbrace{\mathsf{K}^2 \otimes \dots \otimes \mathsf{K}^2}_{j=1} \otimes \mathsf{K}^{-1}\mathsf{F} \otimes \underbrace{\mathsf{K}^{-1} \otimes \dots \otimes \mathsf{K}^{-1}}_{t} \otimes \mathsf{F} \otimes \mathbf{1} \otimes \dots \otimes \mathbf{1}.$$

[2]We note that our convention for the spin-chain representation differs from the one in [23] by the change $\mathfrak{q} \to \mathfrak{q}^{-1}$.

These renormalized powers can also be expressed in terms of the more usual spin-chain operators, and one finds at $p = 2$

$$\Delta^{N-1}(\mathsf{e}) = \mathfrak{q} S^{+(2)} k^2, \qquad \Delta^{N-1}(\mathsf{f}) = \mathfrak{q}^{-1} k^{-2} S^{-(2)},$$

where $\mathfrak{q} = i$ and

$$S^{\pm(2)} = \sum_{1\le j<k\le N-1} \mathfrak{q}^{-\sigma_1^z} \otimes \ldots \otimes \mathfrak{q}^{-\sigma_{j-1}^z} \otimes \sigma_j^{\pm} \otimes 1 \otimes \ldots \otimes 1 \otimes \sigma_k^{\pm} \otimes \mathfrak{q}^{\sigma_{k+1}^z} \otimes \ldots \otimes \mathfrak{q}^{\sigma_N^z}. \tag{151}$$

Appendix B. The proof about the centralizer of $\overline{U}_{\mathfrak{q}} s\ell(2)$ for $\mathfrak{q} = i$

In this section we prove the following theorem.

B.1. **Theorem.** *The centralizer of the $\overline{U}_{\mathfrak{q}} s\ell(2)$ representation at $\mathfrak{q} = i$ – the open XX spin-chain – is generated by*

$$e_k = c_k c_{k+1}^\dagger + c_{k+1} c_k^\dagger + i\big(c_k^\dagger c_k - c_{k+1}^\dagger c_{k+1}\big)$$

and

$$WB_j^+ = (-1)^j \big(c_j^\dagger c_{j+1}^\dagger + i c_j^\dagger c_{j+2}^\dagger - c_{j+1}^\dagger c_{j+2}^\dagger\big),$$
$$WB_j^0 = -\frac{1}{2}(1 + i c_j^\dagger c_{j+1} - c_j^\dagger c_{j+2} + i c_{j+1}^\dagger c_j - 2 c_{j+1}^\dagger c_{j+1} - i c_{j+1}^\dagger c_{j+2} - c_{j+2}^\dagger c_j - i c_{j+2}^\dagger c_{j+1}),$$
$$WB_j^- = (-1)^{j+1} \big(c_j c_{j+1} + i c_j c_{j+2} - c_{j+1} c_{j+2}\big),$$

where $1 \le k \le N-1$ and $1 \le j \le N-2$.

Our proof is computational and consists of few lemmas. Recall first the linear combinations (22) and (31) of the c_j and $c_j^\dagger$ operators – the Fourier transforms $\eta_n^\pm$ of the lattice fermions. We recall also the definition (32) of the zero modes $\eta_0^\pm$ and conjugate to them $\gamma^\mp$ operators. The $\eta_n^\pm$, with $-\frac{N}{2}+1 \le n \le \frac{N}{2}-1$, and $\gamma^\pm$ generate a Clifford algebra of dimension 2^{2N}, see relations in (33). Note that the spin-chain is an irreducible representation of this Clifford algebra. We also recall that each pair of the modes $\eta_n^\pm$, and the pair $\gamma^\pm$, form an $s\ell(2)$ doublet, with respect to the $s\ell(2)$ representation given in (39)-(40), see its action in (41) and (42). Therefore, the $\eta_n^\pm$ and $\gamma^\pm$ change an eigenvalue of the Cartan $S^z = 2\mathsf{h}$ by ± 1. Now, we can formulate and prove the following technical lemma.

B.1.1. **Lemma.** *The centralizer of the $g\ell(1|1)$, i.e., the wall Brauer algebra representation in the XX spin-chain is generated by the fermionic bilinears $\eta_n^+\eta_m^-$, with $-\frac{N}{2}+1 \le n,m \le \frac{N}{2}-1$ and for any $N \in \mathbb{N}$.*

Proof. Define an index set I as the set of integers $-\frac{N}{2}+1 \le n \le \frac{N}{2}-1$. We first note that the centralizer of the algebra generated by S^z – a commutative subalgebra in the $g\ell(1|1)$ representation – has the fermionic bilinears

$$\eta_n^+\eta_m^-, \qquad \text{with } n,m \in I, \tag{152}$$

and

$$\gamma^+\gamma^-, \quad \gamma^+\eta_n^-, \quad \eta_n^+\gamma^-, \qquad \text{with } n \in I, \tag{153}$$

as its generators. Indeed, since the $\eta_n^\pm$ and $\gamma^\pm$ – the generators of the Clifford algebra – change an eigenvalue of S^z by ± 1 it is clear that only monomials in $\eta_n^\pm$ and $\gamma^\pm$ with equal number of '+' and '−' commute with S^z. All such monomials can obviously be generated by the bilinears in (152) and (153), with the use of the (anti)commutation relations in the Clifford algebra.

We then check by direct calculations that any linear combination of the bilinears in (153) does not commute with the fermionic part of $g\ell(1|1)$ – the zero modes $\eta_0^\pm$ – while any linear combination of $\eta_n^+\eta_m^-$ does. It is also straightforward to see that any word from the Clifford algebra involving the bilinears from (153) and any linear combination of such words do not commute with the $\eta_0^\pm$. Finally, recall that the $g\ell(1|1)$ representation in our spin-chain is generated by S^z and $\eta_0^\pm$. We therefore conclude that the centralizer of the $g\ell(1|1)$ action is generated by the $\eta_n^+\eta_m^-$, with $n, m \in I$. □

Consider then a subalgebra $\rho_{g\ell}(\overline{U}_\mathfrak{q} s\ell(2))$ in the (image of) $g\ell(1|1)$, see the definition of $\rho_{g\ell}$ in (10). This subalgebra is generated by $\mathsf{K} = (-1)^{S^z}$ and $\eta_0^\pm$. Our aim is to describe the centralizer of the $\overline{U}_\mathfrak{q} s\ell(2)$ action. Obviously the centralizer of K is generated by all bilinears in the fermionic modes $\eta_n^\pm$ and $\gamma^\pm$. Following the same lines as in the proof of Lem. **B.1.1** we prove our second technical lemma.

B.1.2. **Lemma.** *The centralizer of the $\overline{U}_\mathfrak{q} s\ell(2)$ for $\mathfrak{q} = i$ is generated by the fermionic bilinears $\eta_n^+\eta_m^-$, $\eta_n^+\eta_m^+$, and $\eta_n^-\eta_m^-$, with $-\frac{N}{2}+1 \leq n, m \leq \frac{N}{2}-1$ and for any $N \in \mathbb{N}$.*

Then, note that the centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$ is in the image under the adjoint action of the $s\ell(2)$ algebra on the centralizer of $g\ell(1|1)$, *i.e.*, on the wall Brauer algebra representation. Indeed, in terms of generators for both the centralizers described in Lem. **B.1.1** and Lem. **B.1.2** we have

$$[\mathsf{e}, \eta_n^+\eta_m^-] = \eta_n^+\eta_m^+, \qquad [\mathsf{f}, \eta_n^+\eta_m^-] = \eta_n^-\eta_m^-, \qquad -\frac{N}{2}+1 \leq n, m \leq \frac{N}{2}-1, \tag{154}$$

where we use the fermionic expressions (39) of the e and f generators of the $s\ell(2)$ algebra. We also note that the action of $s\ell(2)$ differentiates the multiplication in the Clifford algebra, *i.e.*, for a and b from the Clifford algebra, we have $\mathsf{e}(ab) = \mathsf{e}(a)b + a\mathsf{e}(b)$, *etc.* Therefore, the equations (154) define the image of any element of the wall Brauer algebra representation under the $s\ell(2)$ action. The centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$ is thus generated by (actually, coincides with) this image of the wall Brauer algebra under the action of $Us\ell(2)$. We finally note that by definition (12)-(15) the lattice W-algebra $\mathcal{W}_{i,N}$ is also generated by the image under the action of $Us\ell(2)$ on the same wall Brauer algebra where just another system of generators, the e_j and WB_j instead of $\eta_n^+\eta_m^-$, is used. The $\mathcal{W}_{i,N}$ is thus isomorphic to the centralizer of $\overline{U}_\mathfrak{q} s\ell(2)$ which finishes the proof of Thm. **B.1**.

Appendix C. Examples of relations in the lattice W-algebra

Here, we give some examples of relations in the lattice W-algebra $\mathcal{W}_{i,N}$.

- 3 **sites:** the defining relations are

$$WB_m^{\pm} WB_m^{\pm} = 0, \tag{155}$$

$$WB_m^0 WB_m^{\pm} = \pm\frac{1}{2} WB_m^{\pm}, \quad WB_m^0 WB_m^0 = \frac{1}{4}(1 - e_m e_{m+1} - e_{m+1} e_m), \tag{156}$$

$$WB_m^+ WB_m^- = WB_m^0 + \frac{1}{2}(1 - e_m e_{m+1} - e_{m+1} e_m), \tag{157}$$

$$e_m WB_m^{\alpha} = WB_m^{\alpha} e_m = e_{m+1} WB_m^{\alpha} = WB_m^{\alpha} e_{m+1} = 0. \tag{158}$$

 These relations allow to construct a basis in the lattice W-algebra on 3 sites. The elements

$$WB_1^{\alpha}, \qquad \alpha = \pm, 0,$$

 together with 5 basis elements from the TL algebra give 8 basis elements in total.

- 4 **sites:** we have the defining relations

$$WB_m^{\pm} WB_{m+1}^{\pm} = 0, \tag{159}$$

$$WB_m^0 WB_{m+1}^0 = \frac{1}{4}(1 - 2e_{2,m} e_{m+1} + 2e_{1,m}), \tag{160}$$

$$WB_{m+1}^0 WB_m^0 = \frac{1}{4}(1 - 2e_{2,m} e_{m+1} - 2e_{1,m+1}), \tag{161}$$

$$WB_m^+ WB_{m+1}^- = WB_{m+1}^0 - e_{m+1} e_m WB_{m+1}^0 + e_{1,m+1} - e_{m+1} e_{2,m} + \frac{1}{2}, \tag{162}$$

$$WB_{m+1}^+ WB_m^- = WB_m^0 + e_{m+1} e_m WB_{m+1}^0 - e_{1,m+1} - e_{m+1} e_{2,m} + \frac{1}{2} \tag{163}$$

 in particular

$$[WB_{m+1}^+, WB_m^-] = WB_m^0 + WB_{m+1}^0 - e_{1,m} - e_{1,m+1}, \tag{164}$$

$$[WB_m^+, WB_{m+1}^-] = WB_m^0 + WB_{m+1}^0 + e_{1,m} + e_{1,m+1}, \tag{165}$$

 and relations with the TL generators

$$e_{m-1} WB_m^{\alpha} e_{m-1} = e_{m+2} WB_m^{\alpha} e_{m+2} = 0,$$

$$[e_m, WB_{m+1}^{\alpha}] = [WB_m^{\alpha}, e_{m+2}],$$

$$e_{1,m} WB_{m+1}^{\alpha} = e_{1,m+1} WB_m^{\alpha}, \qquad WB_{m+1}^{\alpha} e_{1,m} = WB_m^{\alpha} e_{1,m+1}, \quad \alpha = +, -, 0. \tag{166}$$

$$e_{2,m} WB_m^{\alpha} = -\tfrac{1}{2} e_m WB_{m+1}^{\alpha}, \qquad WB_m^{\alpha} e_{2,m} = -\tfrac{1}{2} WB_{m+1}^{\alpha} e_m,$$

$$[e_{m+1}, WB_{1,m}^{\alpha}] = -\tfrac{1}{2} WB_m^{\alpha} + \tfrac{1}{2} WB_{m+1}^{\alpha},$$

 and

$$WB_m^{\pm} e_{m+2} WB_m^{\pm} = WB_{m+1}^{\pm} e_m WB_{m+1}^{\pm} = 0, \tag{167}$$

$$WB_m^0 e_{m+2} WB_m^+ = WB_{m+1}^0 e_m WB_{m+1}^+ = \frac{1}{2}(WB_m^+ e_{m+2} + WB_{m+1}^+ e_m), \tag{168}$$

$$WB^+_m e_{m+2} WB^0_m = WB^+_{m+1} e_m WB^0_{m+1} = -\frac{1}{2}(e_{m+2} WB^+_m + e_m WB^+_{m+1}), \tag{169}$$

$$WB^0_m e_{m+2} WB^0_m = WB^0_{m+1} e_m WB^0_{m+1} = \frac{1}{4}(e_m + e_{m+2} - 2e_{2,m} + 2e_{1,m}e_{m+2} + 2e_m e_{1,m+1} - 2e_m e_{m+1} e_{m+2}), \tag{170}$$

$$WB^+_m e_{m+2} WB^-_m = WB^+_{m+1} e_m WB^-_{m+1} = e_{m+2} WB^0_m + e_m WB^0_{m+1} + e_m e_{m+1} e_{m+2} - \frac{1}{2}(e_m + e_{m+2} - 2e_{2,m} + 2e_{1,m}e_{m+2} + 2e_m e_{1,m+1}) \tag{171}$$

$$WB^-_m e_{m+2} WB^+_m = WB^-_{m+1} e_m WB^+_{m+1} = -WB^0_m e_{m+2} - WB^0_{m+1} e_m - e_m e_{m+1} e_{m+2} + \frac{1}{2}(e_m + e_{m+2} - 2e_{2,m} + 2e_{1,m}e_{m+2} + 2e_m e_{1,m+1}). \tag{172}$$

These relations allow to construct a basis in the lattice W-algebra:

$$WB^\alpha_m, \quad WB^\alpha_{m+1}, \qquad 6 \text{ elements},$$
$$e_{m+2} WB^\alpha_m, \quad WB^\alpha_m e_{m+2}, \quad e_m WB^\alpha_{m+1}, \qquad 9 \text{ elements},$$
$$e_{m+1} e_m WB^\alpha_{m+1}, \qquad 3 \text{ elements},$$

which together with 14 basis elements from the TL algebra give 32 basis elements.

- 5 **sites:** for 5 sites the relations are very bulky and we give only simple examples of them:

$$[WB^+_{m+2}, WB^-_m] = [e_m, [e_{m+1}, WB^0_{m+2}]] - e_{3,m}, \tag{173}$$

$$[WB^+_m, WB^-_{m+2}] = [e_m, [e_{m+1}, WB^0_{m+2}]] + e_{3,m}, \tag{174}$$

$$[WB^0_m, WB^0_{m+2}] = 2e_{3,m}. \tag{175}$$

Appendix D. The triplet W-agebra currents in symplectic fermions

We recall that the triplet W-algebra currents in terms of the symplectic fermions [18]

$$\eta^\pm(z) = \sum z^{-n-1} \eta^\pm_n, \qquad \text{with} \quad \eta^+(z)\eta^-(w) = \frac{1}{(z-w)^2} + \dots, \tag{176}$$

are given by the following expressions

$$T(z) = -:\eta^+(z)\eta^-(z):= \sum z^{-n-2} L_n, \tag{177}$$

$$W^+(z) =: \partial\eta^+(z)\eta^+(z) := \sum z^{-n-3} W^+_n, \tag{178}$$

$$W^0(z) = \frac{1}{2}(:\partial\eta^+(z)\eta^-(z): - :\eta^+(z)\partial\eta^-(z):) = \sum z^{-n-3} W^0_n, \tag{179}$$

$$W^-(z) =: \partial\eta^-(z)\eta^-(z) := \sum z^{-n-3} W^-_n. \tag{180}$$

We also introduce modes of some composite currents

$$:T(z)T(z):= \sum z^{-n-4} (L^2)_n, \tag{181}$$

$$:T(z)W^\alpha(z):= \sum z^{-n-5} (LW^\alpha)_n. \tag{182}$$

These modes are expressed in the modes of symplectic fermions

$$L_n = -\sum_{j_1+j_2=n} :\eta^+_{j_1}\eta^-_{j_2}:, \tag{183}$$

$$(L^2)_n = -\frac{1}{2}\sum_{j_1+j_2=n} ((j_1+1)(j_1+2)+(j_2+1)(j_2+2)) :\eta^+_{j_1}\eta^-_{j_2}:, \tag{184}$$

$$W^+_n = -\sum_{j_1+j_2=n} (j_1+1)\eta^+_{j_1}\eta^+_{j_2}, \tag{185}$$

$$W^0_n = -\sum_{j_1+j_2=n} (j_1-j_2)\eta^+_{j_1}\eta^-_{j_2}, \tag{186}$$

$$W^+_n = -\sum_{j_1+j_2=n} (j_1+1)\eta^-_{j_1}\eta^-_{j_2}, \tag{187}$$

$$(LW^+)_n = \sum_{j_1+j_2=n} (-\frac{1}{3}(j_2+1)(j_2+2)(j_2+3)+\frac{1}{2}(j_1+1)(j_2+1)(j_2+2))\eta^+_{j_1}\eta^+_{j_2}. \tag{188}$$

The modes of these currents satisfy the commutation relations [11]

$$[L_n, L_m] = (n-m)L_{n+m} - \frac{1}{6}n(n^2-1)\delta_{n+m,0}, \tag{189}$$

$$[L_n, W^\alpha_m] = (2n-m)W^\alpha_{n+m}, \tag{190}$$

$$\begin{aligned}[W^\alpha_n, W^\beta_m] = g^{\alpha\beta}\Big(2(n-m)(L^2)_{n+m} - \frac{1}{4}(n-m)(2n+m+4)(2m+n+4)L_{n+m} - \\ - \frac{1}{120}n(n^2-1)(n^2-4)\delta_{n+m,0}\Big) + \\ + f^{\alpha\beta}_\gamma\Big(\frac{12}{5}(LW^\gamma)_{n+m} + \frac{1}{10}(2n^2+2m^2-21nm-36n-36m-76)W^\gamma_{n+m}\Big).\end{aligned} \tag{191}$$

where $g^{\alpha\beta}$ and $f^{\alpha\beta}_\gamma$ are defined after (1).

Our aim in the paper is to show that these commutation relations can be obtained from the lattice W-algebra $\mathcal{W}_{i,N}$.

Appendix E. Commutation relations in the $\mathcal{W}_{i,N}$ algebra

We collect here a list of commutators of the Fourier modes in $\mathcal{W}_{i,N}$.

$$[H^0_n, W^{\alpha,0}_m] = 2\sin\pi\frac{2n-m}{2N}W^{\alpha,1}_{n+m} + 2\sin\pi\frac{2n+m}{2N}W^{\alpha,1}_{n-m}, \tag{192}$$

$$\begin{aligned}[H^0_n, W^{\alpha,1}_m] = 2\sin\pi\frac{3n-m}{2N}W^{\alpha,2}_{n+m} - 2\sin\pi\frac{3n+m}{2N}W^{\alpha,2}_{n-m} - \\ - \sin\pi\frac{n}{N}\big(\cos\pi\frac{n-m}{2N}W^{\alpha,0}_{n+m} - \cos\pi\frac{n+m}{2N}W^{\alpha,0}_{n-m}\big),\end{aligned} \tag{193}$$

$$\begin{aligned}[H^1_n, W^{\alpha,0}_m] = 2\sin\pi\frac{n-m}{N}W^{\alpha,2}_{n+m} + 2\sin\pi\frac{n+m}{N}W^{\alpha,2}_{n-m} + \\ + 2\cos\pi\frac{n}{2N}\sin\pi\frac{m}{2N}\big(\cos\pi\frac{n-m}{2N}W^{\alpha,0}_{n+m} - \cos\pi\frac{n+m}{2N}W^{\alpha,0}_{n-m}\big),\end{aligned} \tag{194}$$

$$[H^1_n, W^{\alpha,1}_m] = 2\sin\pi\frac{3n-2m}{2N}W^{\alpha,3}_{n+m} - 2\sin\pi\frac{3n+2m}{2N}W^{\alpha,3}_{n-m}+ \tag{195}$$
$$+\big(\cos\pi\frac{n}{N}\sin\pi\frac{n}{2N} - \sin\pi\frac{3n-2m}{2N}\big)W^{\alpha,1}_{n+m}-$$
$$-\big(\cos\pi\frac{n}{N}\sin\pi\frac{n}{2N} - \sin\pi\frac{3n+2m}{2N}\big)W^{\alpha,1}_{n-m},$$

(196)
$$[W^{\alpha,0}_n, W^{\beta,0}_m] = f^{\alpha\beta}_\gamma\Big(2\cos\pi\frac{n-m}{N}W^{\gamma,2}_{n+m} + 2\cos\pi\frac{n+m}{N}W^{\gamma,2}_{n-m}-$$
$$-\frac{1}{2}\big(\cos\pi\frac{n}{N}+\cos\pi\frac{m}{N}+\cos\pi\frac{n-m}{N}+1\big)W^{\gamma,0}_{n+m}-$$
$$-\frac{1}{2}\big(\cos\pi\frac{n}{N}+\cos\pi\frac{m}{N}+\cos\pi\frac{n+m}{N}+1\big)W^{\gamma,0}_{n-m}\Big)+$$
$$+ g^{\alpha\beta}\Big(4\sin\pi\frac{n-m}{N}H^3_{n+m} + 4\sin\pi\frac{n+m}{N}H^3_{n-m}-$$
$$-\big(\sin\pi\frac{n}{N}-\sin\pi\frac{m}{N}+3\sin\pi\frac{n-m}{N}\big)H^1_{n+m}-$$
$$-\big(\sin\pi\frac{n}{N}+\sin\pi\frac{m}{N}+3\sin\pi\frac{n-m}{N}\big)H^1_{n-m}\Big),$$

(197)
$$[W^{\alpha,0}_n, W^{\beta,1}_m] = f^{\alpha\beta}_\gamma\Big(2\cos\pi\frac{3n-2m}{2N}W^{\gamma,3}_{n+m} - 2\cos\pi\frac{3n+2m}{2N}W^{\gamma,3}_{n-m}-$$
$$-\big(\cos\pi\frac{n}{2N}\cos\pi\frac{n}{N}+\cos\pi\frac{3n-2m}{2N}\big)W^{\gamma,1}_{n+m}+$$
$$+\big(\cos\pi\frac{n}{2N}\cos\pi\frac{n}{N}+\cos\pi\frac{3n+2m}{2N}\big)W^{\gamma,1}_{n-m}\Big)+$$
$$+ g^{\alpha\beta}\Big(4\sin\pi\frac{3n-2m}{2N}H^4_{n+m} - 4\sin\pi\frac{3n+2m}{2N}H^4_{n-m}-$$
$$-2\big(\cos\pi\frac{n}{2N}\sin\pi\frac{n}{N}+2\sin\pi\frac{3n-2m}{2N}\big)H^2_{n+m}+$$
$$+2\big(\cos\pi\frac{n}{2N}\sin\pi\frac{n}{N}+2\sin\pi\frac{3n+2m}{2N}\big)H^2_{n-m}+$$
$$+2\sin\pi\frac{n}{N}\cos\pi\frac{m}{2N}\cos\pi\frac{n-m}{2N}H^0_{n+m}-$$
$$-2\sin\pi\frac{n}{N}\cos\pi\frac{m}{2N}\cos\pi\frac{n+m}{2N}H^0_{n-m}\Big),$$

(198)
$$[W^{\alpha,1}_n, W^{\beta,0}_m] = f^{\alpha\beta}_\gamma\Big(2\cos\pi\frac{2n-3m}{2N}W^{\gamma,3}_{n+m} + 2\cos\pi\frac{2n+3m}{2N}W^{\gamma,3}_{n-m}-$$
$$-\big(\cos\pi\frac{m}{2N}\cos\pi\frac{m}{N}+\cos\pi\frac{2n-3m}{2N}\big)W^{\gamma,1}_{n+m}+$$
$$-\big(\cos\pi\frac{m}{2N}\cos\pi\frac{m}{N}+\cos\pi\frac{2n+3m}{2N}\big)W^{\gamma,1}_{n-m}\Big)+$$

$$
\begin{aligned}
&+ g^{\alpha\beta}\Big(4\sin\pi\frac{2n-3m}{2N}H^4_{n+m} + 4\sin\pi\frac{2n+3m}{2N}H^4_{n-m} + \\
&\quad + 2\big(\cos\pi\frac{m}{2N}\sin\pi\frac{m}{N} - 2\sin\pi\frac{2n-3m}{2N}\big)H^2_{n+m} - \\
&\quad - 2\big(\cos\pi\frac{m}{2N}\sin\pi\frac{m}{N} + 2\sin\pi\frac{2n+3m}{2N}\big)H^2_{n-m} - \\
&\quad - 2\sin\pi\frac{m}{N}\cos\pi\frac{n}{2N}\cos\pi\frac{n-m}{2N}H^0_{n+m} + \\
&\quad + 2\sin\pi\frac{m}{N}\cos\pi\frac{n}{2N}\cos\pi\frac{n+m}{2N}H^0_{n-m}\Big),
\end{aligned}
$$

(199)

$$
\begin{aligned}
[W^{\alpha,1}_n, W^{\beta,1}_m] &= f^{\alpha\beta}_\gamma\Big(2\cos\pi\frac{3n-3m}{2N}W^{\gamma,4}_{n+m} - 2\cos\pi\frac{3n+3m}{2N}W^{\gamma,4}_{n-m} - \\
&\quad - \frac{1}{2}\big(\cos\pi\frac{n-m}{2N} + 3\cos\pi\frac{3n-3m}{2N}\big)W^{\gamma,2}_{n+m} + \\
&\quad + \frac{1}{2}\big(\cos\pi\frac{n+m}{2N} + 3\cos\pi\frac{3n+2m}{2N}\big)W^{\gamma,2}_{n-m} + \\
&\quad + \frac{1}{2}\cos\pi\frac{n-m}{2N}\sin\pi\frac{n}{N}\sin\pi\frac{m}{N}W^{\gamma,0}_{n+m} + \\
&\quad + \frac{1}{2}\cos\pi\frac{n+m}{2N}\sin\pi\frac{n}{N}\sin\pi\frac{m}{N}W^{\gamma,0}_{n-m}\Big) \\
&+ g^{\alpha\beta}\Big(4\sin\pi\frac{3n-3m}{2N}H^5_{n+m} - 4\sin\pi\frac{3n+3m}{2N}H^5_{n-m} - \\
&\quad - \big(\sin\pi\frac{n-m}{2N} + 5\sin\pi\frac{3n-3m}{2N}\big)H^3_{n+m} + \\
&\quad + \big(\sin\pi\frac{n+m}{2N} + 5\sin\pi\frac{3n+3m}{2N}\big)H^3_{n-m} - \\
&\quad - \frac{1}{4}\big(\cos\pi\frac{n+m}{N}\sin\pi\frac{n-m}{2N} - 3\sin\pi\frac{n-m}{2N} - 5\sin\pi\frac{3n-3m}{2N}\big)H^1_{n+m} + \\
&\quad + \frac{1}{4}\big(\cos\pi\frac{n-m}{N}\sin\pi\frac{n+m}{2N} - 3\sin\pi\frac{n+m}{2N} - 5\sin\pi\frac{3n+3m}{2N}\big)H^1_{n-m}\Big).
\end{aligned}
$$

References

[1] E. Date, M. Jimbo, T. Miwa and M. Okado, *Automorphic properties of local height probabilities for integrable solid-on-solid models*, Phys. Rev. B35 (1987) 2105.

[2] N. Read and H. Saleur, *Enlarged symmetry algebras of spin chains, loop models, and S-matrices*, Nucl. Phys. B **777**, 263 (2007).

[3] N. Read and H. Saleur, *Associative-algebraic approach to logarithmic conformal field theories*, Nucl. Phys. B **777**, 316 (2007).

[4] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I.Yu. Tipunin, *Logarithmic extensions of minimal models: characters and modular transformations*, Nucl. Phys. B757 (2006) 303-343, hep-th/0606196.

[5] A. Gaynutdinov, N. Read and H. Saleur, *Continuum limit and symmetries of the periodic gl(1/1) spin chain*, arXiv:1112.3403

[6] A. Gaynutdinov, N. Read and H. Saleur, *Bimodule structure in the periodic gl(1/1) spin chain*, arXiv:1112.3407

[7] P. Pearce, J. Rasmussen and J.B. Zuber, *Logarithmic Minimal Models*, J. Stat. Mech. (2006) 0611:017.
[8] J. Rasmussen and P. Pearce, *Fusion Algebras of Logarithmic Minimal Models*, J. Phys. A40 (2007) 13711-13734.
[9] N. Read and H. Saleur, *Exact spectra of conformal supersymmetric nonlinear sigma models in two dimensions*, Nucl. Phys. B613 (2001) 409.
[10] W.M. Koo and H. Saleur, *Representations of the Virasoro algebra from lattice models*, Int. J. of Mod. Phys. A8 (1993) 5165.
[11] M.R. Gaberdiel, H.G. Kausch, *A Rational Logarithmic Conformal Field Theory*, Phys.Lett. B386 (1996) 131-137, hep-th/9606050
[12] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I.Yu. Tipunin, *Kazhdan–Lusztig correspondence for the representation category of the triplet W-algebra in logarithmic CFT*, Theor. Math. Phys. 148 (2006) 1210–1235.
[13] Y.-Z. Huang, J. Lepowsky, and L. Zhang, *Logarithmic tensor category theory for generalized modules for a conformal vertex algebra, I-VI*, arXiv:1012.4193; arXiv:1012.4196; arXiv:1012.4197; arXiv:1012.4198; arXiv:1012.4199; arXiv:1012.4202.
[14] D. Adamović, A. Milas, *On W-algebras associated to (2,p) minimal models and their representations*, arXiv:0908.4053
[15] M. R. Gaberdiel, I. Runkel, J. Phys. A41 (2008) 075402.
[16] A.M. Gainutdinov, Theor. Math. Phys., 159(2): 575-586 (2009).
[17] J. Fuchs, Ch. Schweigert, C. Stigner, *Higher genus mapping class group invariants from factorizable Hopf algebras*, ZMP-HH/12-13, arXiv:1207.6863.
[18] H.G. Kausch, *Curiosities at c=-2*, hep-th/9510149
[19] J. Rasmussen, *W-Extended Logarithmic Minimal Models*, arXiv:0805.2991
[20] C. Candu, V. Mitev, T. Quella, H. Saleur, V. Schomerus, *The Sigma Model on Complex Projective Superspaces*, JHEP 1002:015,2010
[21] P.P. Martin and D. McAnally, *On Commutants, Dual Pairs and Non-Semisimple Algebras from Statistical Mechanics*, Int. J. Mod. Phys. A 7, Supp. 1B, 675 (1992); P.P. Martin, *On Schur–Weyl duality, A_n Hecke algebras and quantum $sl(N)$ on $\otimes^{n+1}\mathbb{C}^N$*, Int. J. Mod. Phys. A 7, Supp. 1B, 645 (1992).
[22] S. Donkin, *The q-Schur algebra*, London Mathematical Society Lecture Note Series, vol. 253, Cambridge Univ. Press, Cambridge,1999, 179pp. ISBN 0-521-64558-1
[23] V. Pasquier and H. Saleur, Nucl. Phys. B330 (1990) 523.
[24] H. Araki, Comm. Math. PHys. 132 (1990) 155.
[25] P.V. Bushlanov, B.L. Feigin, A.M. Gainutdinov, I.Yu. Tipunin, Nucl. Phys. B 818(3), 179-195 (2009)
[26] B.L. Feigin, A.M. Gainutdinov, A.M. Semikhatov, and I.Yu. Tipunin, *Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center*, Commun. Math. Phys. 265 (2006) 47–93. [hep-th/0504093].
[27] K. Nagatomo and A. Tsuchiya, *The triplet vertex-operator algebra $W(p)$ and the restricted quantum group at root of unity*, arXiv:0902.4607 [math.QA].
[28] P.V. Bushlanov, A.M. Gainutdinov, and I.Yu. Tipunin, *Kazhdan-Lusztig equivalence and fusion of Kac modules in Virasoro logarithmic models*, arXiv:1102.0271.
[29] A. Sergeev, *An analog of the classical invariant theory for Lie superalgebras I, II*, Michigan Math. J. 49, 113, 147 (2001).
[30] T. Deguchi, K. Fabricius and B. Mc Coy, *The $sl(2)$ loop algebra symmetry of the six-vertex model at roots of unity*, J. Stat. Phys. 102 (2001) 701.
[31] C. Ahn and K. Shigemoto: Modern Phys. Lett. A 6 (1991) 3509.

[32] A.M. Gainutdinov and R. Vasseur, *Lattice fusion rules and logarithmic operator product expansions*, arXiv:1203.6289
[33] R.J. Baxter, *Exactly Solved Models in Statistical Mechanics*.
[34] V. Kac, *Infinite-dimensional Lie algebras*, Cambridge (1994) 424pp.
[35] E. Date and S. Roan, *The structure of quotients of the Onsager algebra by closed ideals*, J. Phys. A: Math. Gen. 33 (2000) 3275.
[36] M. Gaberdiel, I. Runkel and S. Wood, *Fusion rules and boundary conditions in the $c = 0$ triplet model*, J. Phys. A: Math. Theor. **42**, 325403 (2009).
[37] A.M. Gainutdinov, I.Yu. Tipunin, in preparation.

AMG: Institut de Physique Théorique, CEA Saclay, Gif Sur Yvette, 91191, France

E-mail address: azat.gaynutdinov@cea.fr

HS: Institut de Physique Théorique, CEA Saclay, Gif Sur Yvette, 91191, France.
Department of Physics and Astronomy, University of Southern California, Los Angeles, CA 90089, USA

E-mail address: hubert.saleur@cea.fr

IYuT: Tamm Theory Division, Lebedev Physics Institute, Leninski pr., 53, Moscow, Russia, 119991

E-mail address: tipunin@gmail.com